



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> C12N 15/82, 15/31, A01N 63/02	<b>A3</b>	<b>(11) International Publication Number:</b> WO 00/28055 <b>(43) International Publication Date:</b> 18 May 2000 (18.05.00)
<b>(21) International Application Number:</b> PCT/US99/26039 <b>(22) International Filing Date:</b> 4 November 1999 (04.11.99) <b>(30) Priority Data:</b> 60/107,243 5 November 1998 (05.11.98) US <b>(71) Applicant:</b> EDEN BIOSCIENCE CORPORATION [US/US]; 11816 North Creek Parkway N., Bothell, WA 98011-8205 (US). <b>(72) Inventors:</b> WEI, Zhong-Min; 8230 125th Court, Kirkland, WA 98034 (US). SCHADING, Richard, L.; 3607 Carriage Gate Drive, West Melbourne, FL 32904 (US). <b>(74) Agents:</b> GOLDMAN, Michael, L. et al.; Nixon Peabody LLP, Clinton Square, P.O. Box 1051, Rochester, NY 14603 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>  <b>(88) Date of publication of the international search report:</b> 14 September 2000 (14.09.00)
<b>(54) Title:</b> HYPERSENSITIVE RESPONSE ELICITOR-INDUCED STRESS RESISTANCE  <b>(57) Abstract</b>  The present invention is directed to imparting stress resistance to plants. This can be achieved by applying a hypersensitive response elicitor in a non-infectious form to plants or plant seeds under conditions effective to impart stress resistance to plants or plants grown from the plant seeds. Alternatively, transgenic plants or plant seeds transformed with a DNA molecule encoding the elicitor can be provided and the transgenic plants or plants resulting from the transgenic plant seeds are grown under conditions effective to impart stress resistance to plants or plants grown from the plant seeds.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

SEP 22 2000

## **HYPERSENSITIVE RESPONSE ELICITOR-INDUCED STRESS RESISTANCE**

5

### **FIELD OF THE INVENTION**

10           The present invention relates to imparting stress resistance to plants with a hypersensitive response elicitor.

### **BACKGROUND OF THE INVENTION**

15           Under both natural and agricultural conditions, plants are exposed to various forms of environmental stress. Stress is mainly measured with respect to growth (i.e. biomass accumulation) or with respect to the primary assimilation processes (i.e. carbon dioxide and mineral intake). Soil water deficits, suboptimal and supraoptimal temperatures, salinity, and poor aeration of soils may each cause some  
20   growth restrictions during the growing season, so that the yield of plants at the end of the season expresses only a small fraction of their genetic potential. Indeed, it is estimated that in the United States the yield of field-grown crops is only 22% of genetic potential. The same physicochemical factors can become extreme in some habitats, such as deserts or marshes, and only specially adapted vegetation can  
25   complete its life cycle in the unusually hostile conditions. In less extreme environments, individual plants can become acclimated to changes in water potential, temperature, salinity, and oxygen deficiency so that their fitness for those environments improves. Some species are better able to adapt than others, and various anatomical, structural, and biochemical mechanisms account for acclimation.

30           Under natural and agriculture conditions, plants must constantly endure stress. Some environmental factors can become stressful in a very short period of time (e.g., high or low temperature) or may take long periods of time to stress plants (e.g., soil water content or mineral nutrients). Generally, environmental stress effecting plants can be in the form of climate related stress, air pollution stress,

chemical stress, and nutritional stress. Examples of climate related stress include drought, water, frost, cold temperature, high temperature, excessive light, and insufficient light. Air pollution stress can be in the form of carbon dioxide, carbon monoxide, sulfur dioxide, NO<sub>x</sub>, hydrocarbons, ozone, ultraviolet radiation, and acidic rain. Chemical stress can result from application of insecticides, fungicides, herbicides, and heavy metals. Nutritional stress can be caused by fertilizers, micronutrients, and macronutrients.

For most plants, water is essential for growth. Some plants are able to preserve some water in the soil for later use, while others complete their life cycles during a wet season before the onset of any drought. Other plants are able to aggressively consume water to save themselves while causing water deprivation for other plants in that location. Plants lacking any of these capabilities are severely hampered by the absence of water.

Chilling injury occurs in sensitive species at temperatures that are too low for normal growth but not sufficiently low to form ice. Such injury typically occurs in species of tropical or subtropical origin. When chilling occurs, discoloration or lesions appear on leaves giving them a water-soaked appearance. If roots are chilled, the plants may wilt. On the other hand, freezing temperatures and the accompanying formation of ice crystals in plants can be lethal if ice crystals extend into protoplasts or remain for long periods.

Stress is also caused by the other temperature extremes with few plants being able to survive high temperatures. When higher plant cells or tissues are dehydrated or are not growing, they can survive higher temperatures than cells which are hydrated, vegetative, and growing. Tissues which are actively growing can rarely survive at temperatures above 45°C.

High salt concentrations are another form of environmental stress which can afflict plants. In natural conditions, such high concentrations of salt are found close to seashores and estuaries. Farther inland, natural salt may seep from geological deposits adjoining agricultural areas. In addition, salt can accumulate in irrigation water when pure water is evaporated or transpired from soil. About 10% of all irrigated farmland is effected by high salt concentrations. High salt content not

only injures plants but degrades soil structure by decreasing porosity and water permeability.

Air pollution in the form of ozone, carbon dioxide, carbon monoxide, sulfur dioxide, NO<sub>x</sub>, and hydrocarbons can very adversely effect plant growth by creating smog and environmental warming.

The present invention is directed to overcoming various forms of environmental stress and imparting resistance in plants to such stress.

### SUMMARY OF THE INVENTION

10

The present invention relates to the use of a hypersensitive response elicitor protein or polypeptide to impart stress resistance to plants. In one embodiment of the present invention, the hypersensitive response elicitor protein or polypeptide is applied to plants or plant seeds under conditions effective to impart stress resistance. Alternatively, stress resistance is imparted by providing a transgenic plant or plant seed transformed with a DNA molecule which encodes for a hypersensitive response elicitor protein or polypeptide and growing the transgenic plant or plants produced from the transgenic plant seeds under conditions effective to impart stress resistance.

20

Stress encompasses any environmental factor having an adverse effect on plant physiology and development. Examples of such environmental stress include climate-related stress (e.g., drought, water, frost, cold temperature, high temperature, excessive light, and insufficient light), air pollution stress (e.g., carbon dioxide, carbon monoxide, sulfur dioxide, NO<sub>x</sub>, hydrocarbons, ozone, ultraviolet radiation, acidic rain), chemical (e.g., insecticides, fungicides, herbicides, heavy metals), and nutritional stress (e.g., fertilizer, micronutrients, macronutrients). Applicants have found that use of hypersensitive response elicitors in accordance with the present invention impart resistance to plants against such forms of environmental stress.

30

### DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the use of a hypersensitive response elicitor protein or polypeptide to impart stress resistance to plants. In one

embodiment of the present invention, the hypersensitive response elicitor protein or polypeptide is applied to plants or plant seeds under conditions effective to impart stress resistance. Alternatively, the stress resistance is imparted by providing a transgenic plant or plant seed transformed with a DNA molecule which encodes for a hypersensitive response elicitor protein or polypeptide and growing the transgenic plant or plants produced from the transgenic plant seeds under conditions effective to impart stress resistance.

The hypersensitive response elicitor polypeptides or proteins according to the present invention are derived from hypersensitive response elicitor polypeptides or proteins of a wide variety of fungal and bacterial pathogens. Such polypeptides or proteins are able to elicit local necrosis in plant tissue contacted by the elicitor. Examples of suitable bacterial sources of polypeptide or protein elicitors include *Erwinia*, *Pseudomonas*, and *Xanthomonas* species (e.g., the following bacteria: *Erwinia amylovora*, *Erwinia chrysanthemi*, *Erwinia stewartii*, *Erwinia carotovora*, *Pseudomonas syringae*, *Pseudomonas solanacearum*, *Xanthomonas campestris*, and mixtures thereof). In addition to hypersensitive response elicitors from these Gram negative bacteria, it is possible to use elicitors from Gram positive bacteria. One example is *Clavibacter michiganensis* subsp. *sepedonicus*.

An example of a fungal source of a hypersensitive response elicitor protein or polypeptide is *Phytophthora*. Suitable species of *Phytophthora* include *Phytophthora parasitica*, *Phytophthora cryptogea*, *Phytophthora cinnamomi*, *Phytophthora capsici*, *Phytophthora megasperma*, and *Phytophthora citrophthora*.

The hypersensitive response elicitor polypeptide or protein from *Erwinia chrysanthemi* has an amino acid sequence corresponding to SEQ. ID. No. 1 as follows:

Met	Gln	Ile	Thr	Ile	Lys	Ala	His	Ile	Gly	Gly	Asp	Leu	Gly	Val	Ser	1	5	10	15
Gly	Leu	Gly	Ala	Gln	Gly	Leu	Lys	Gly	Leu	Asn	Ser	Ala	Ala	Ser	Ser	20	25	30	
Leu	Gly	Ser	Ser	Val	Asp	Lys	Leu	Ser	Ser	Thr	Ile	Asp	Lys	Leu	Thr	35	40	45	
Ser	Ala	Leu	Thr	Ser	Met	Met	Phe	Gly	Gly	Ala	Leu	Ala	Gln	Gly	Leu	50	55	60	

- 5 -

	Gly	Ala	Ser	Ser	Lys	Gly	Leu	Gly	Met	Ser	Asn	Gln	Leu	Gly	Gln	Ser	
	65					70					75					80	
	Phe	Gly	Asn	Gly	Ala	Gln	Gly	Ala	Ser	Asn	Leu	Leu	Ser	Val	Pro	Lys	
					85					90					95		
5	Ser	Gly	Gly	Asp	Ala	Leu	Ser	Lys	Met	Phe	Asp	Lys	Ala	Leu	Asp	Asp	
				100					105					110			
	Leu	Leu	Gly	His	Asp	Thr	Val	Thr	Lys	Leu	Thr	Asn	Gln	Ser	Asn	Gln	
			115					120					125				
10	Leu	Ala	Asn	Ser	Met	Leu	Asn	Ala	Ser	Gln	Met	Thr	Gln	Gly	Asn	Met	
		130					135					140					
	Asn	Ala	Phe	Gly	Ser	Gly	Val	Asn	Asn	Ala	Leu	Ser	Ser	Ile	Leu	Gly	
	145					150					155					160	
	Asn	Gly	Leu	Gly	Gln	Ser	Met	Ser	Gly	Phe	Ser	Gln	Pro	Ser	Leu	Gly	
					165					170					175		
15	Ala	Gly	Gly	Leu	Gln	Gly	Leu	Ser	Gly	Ala	Gly	Ala	Phe	Asn	Gln	Leu	
				180					185					190			
	Gly	Asn	Ala	Ile	Gly	Met	Gly	Val	Gly	Gln	Asn	Ala	Ala	Leu	Ser	Ala	
			195					200					205				
20	Leu	Ser	Asn	Val	Ser	Thr	His	Val	Asp	Gly	Asn	Asn	Arg	His	Phe	Val	
		210					215					220					
	Asp	Lys	Glu	Asp	Arg	Gly	Met	Ala	Lys	Glu	Ile	Gly	Gln	Phe	Met	Asp	
	225					230					235					240	
	Gln	Tyr	Pro	Glu	Ile	Phe	Gly	Lys	Pro	Glu	Tyr	Gln	Lys	Asp	Gly	Trp	
				245						250					255		
25	Ser	Ser	Pro	Lys	Thr	Asp	Asp	Lys	Ser	Trp	Ala	Lys	Ala	Leu	Ser	Lys	
				260					265					270			
	Pro	Asp	Asp	Asp	Gly	Met	Thr	Gly	Ala	Ser	Met	Asp	Lys	Phe	Arg	Gln	
			275					280					285				
30	Ala	Met	Gly	Met	Ile	Lys	Ser	Ala	Val	Ala	Gly	Asp	Thr	Gly	Asn	Thr	
		290				295						300					
	Asn	Leu	Asn	Leu	Arg	Gly	Ala	Gly	Gly	Ala	Ser	Leu	Gly	Ile	Asp	Ala	
	305					310					315					320	
	Ala	Val	Val	Gly	Asp	Lys	Ile	Ala	Asn	Met	Ser	Leu	Gly	Lys	Leu	Ala	
					325					330					335		
35	Asn	Ala															

This hypersensitive response elicitor polypeptide or protein has a molecular weight of 34 kDa, is heat stable, has a glycine content of greater than 16%, and contains

substantially no cysteine. The *Erwinia chrysanthemi* hypersensitive response elicitor polypeptide or protein is encoded by a DNA molecule having a nucleotide sequence corresponding to SEQ. ID. No. 2 as follows:

5	CGATTTTACC CGGGTGAACG TGCTATGACC GACAGCATCA CGGTATTCTGA CACCGTTACG	60
	GCGTTTATGG CCGCGATGAA CCGGCATCAG GCGGCGCGCT GGTCGCCGCA ATCCGGCGTC	120
	GATCTGGTAT TTCAGTTTGG GGACACCGGG CGTGAACCTCA TGATGCAGAT TCAGCCGGGG	180
	CAGCAATATC CCGGCATGTT GCGCACGCTG CTCGCTCGTC GTTATCAGCA GGCGGCAGAG	240
	TGCGATGGCT GCCATCTGTG CCTGAACGGC AGCGATGTAT TGATCCTCTG GTGGCCGCTG	300
10	CCGTCGGATC CCGGCAGTTA TCCGCAGGTG ATCGAACGTT TGTTTGAACCT GGCGGGAATG	360
	ACGTTGCCGT CGCTATCCAT AGCACCGACG GCGCGTCCGC AGACAGGGAA CGGACGCGCC	420
	CGATCATTA GATAAAGGCG GCTTTTTTTT TTGCAAAACG GTAACGGTGA GGAACCGTTT	480
	CACCGTCGGC GTCACCTCAGT AACAACTATC CATCATGATG CCTACATCGG GATCGGCGTG	540
	GGCATCCGTT GCAGATACTT TTGCGAACAC CTGACATGAA TGAGGAAACG AAATTATGCA	600
15	AATTACGATC AAAGCGCACA TCGGCGGTGA TTTGGGCGTC TCCGGTCTGG GGCTGGGTGC	660
	TCAGGGACTG AAAGGACTGA ATTCCGCGGC TTCATCGCTG GGTTCAGCG TGGATAAACT	720
	GAGCAGCACC ATCGATAAGT TGACCTCCGC GCTGACTTCG ATGATGTTTG GCGGCGCGCT	780
	GGCGCAGGGG CTGGGCGCCA GCTCGAAGGG GCTGGGGATG AGCAATCAAC TGGGCCAGTC	840
	TTTCGGCAAT GGCGCGCAGG GTGCGAGCAA CCTGCTATCC GTACCGAAAT CCGGCGGCGA	900
20	TGCGTTGTCA AAAATGTTTG ATAAAGCGCT GGACGATCTG CTGGGTCATG ACACCGTGAC	960
	CAAGCTGACT AACCAGAGCA ACCAACTGGC TAATTCAATG CTGAACGCCA GCCAGATGAC	1020
	CCAGGGTAAT ATGAATGCGT TCGGCAGCGG TGTGAACAAC GCACTGTCTG CCATTCTCGG	1080
	CAACGGTCTC GGCCAGTCGA TGAGTGGCTT CTCTCAGCCT TCTCTGGGGG CAGGCGGCTT	1140
	GCAGGGCCTG AGCGGCGCGG GTGCATTCAA CCAGTTGGGT AATGCCATCG GCATGGGCGT	1200
25	GGGGCAGAAT GCTGCGCTGA GTGCGTTGAG TAACGTCAGC ACCCACGTAG ACGGTAACAA	1260
	CCGCCACTTT GTAGATAAAG AAGATCGCGG CATGGCGAAA GAGATCGGCC AGTTTATGGA	1320
	TCAGTATCCG GAAATATTCG GTAAACCGGA ATACCAGAAA GATGGCTGGA GTTCGCCGAA	1380
	GACGGACGAC AAATCCTGGG CTAAAGCGCT GAGTAAACCG GATGATGACG GTATGACCGG	1440
	CGCCAGCATG GACAAATTCC GTCAGGCGAT GGGTATGATC AAAAGCGCGG TGCGGGGTGA	1500
30	TACCGGCAAT ACCAACCTGA ACCTGCGTGG CGCGGGCGGT GCATCGCTGG GTATCGATGC	1560
	GGCTGTCTGC GCGGATAAAA TAGCCAACAT GTCGCTGGGT AAGCTGGCCA ACGCCTGATA	1620



- 7 -

ATCTGTGCTG GCCTGATAAA GCGGAAACGA AAAAAGAGAC GGGGAAGCCT GTCTCTTTTC 1680  
 TTATTATGCG GTTTATGCGG TTACCTGGAC CGGTTAATCA TCGTCATCGA TCTGGTACAA 1740  
 ACGCACATTT TCCCGTTCAT TCGCGTCGTT ACGCGCCACA ATCGCGATGG CATCTTCCTC 1800  
 GTCGCTCAGA TTGCGCGGCT GATGGGGAAC GCCGGGTGGA ATATAGAGAA ACTCGCCGGC 1860  
 5 CAGATGGAGA CACGTCTGCG ATAAATCTGT GCCGTAACGT GTTTCTATCC GCCCCTTTAG 1920  
 CAGATAGATT GCGGTTTCGT AATCAACATG GTAATGCGGT TCCGCCTGTG CGCCGGCCGG 1980  
 GATCACCACA ATATTCATAG AAAGCTGTCT TGCACCTACC GTATCGCGGG AGATACCGAC 2040  
 AAAATAGGGC AGTTTTTGCG TGGTATCCGT GGGGTGTTCC GGCCTGACAA TCTTGAGTTG 2100  
 GTTCGTCATC ATCTTTCTCC ATCTGGGCGA CCTGATCGGT T 2141  
 10

The hypersensitive response elicitor polypeptide or protein derived  
 from *Erwinia amylovora* has an amino acid sequence corresponding to SEQ. ID.  
 No. 3 as follows:

15 Met Ser Leu Asn Thr Ser Gly Leu Gly Ala Ser Thr Met Gln Ile Ser  
 1 5 10 15  
 Ile Gly Gly Ala Gly Gly Asn Asn Gly Leu Leu Gly Thr Ser Arg Gln  
 20 20 25 30  
 Asn Ala Gly Leu Gly Gly Asn Ser Ala Leu Gly Leu Gly Gly Gly Asn  
 35 40 45  
 Gln Asn Asp Thr Val Asn Gln Leu Ala Gly Leu Leu Thr Gly Met Met  
 50 55 60  
 Met Met Met Ser Met Met Gly Gly Gly Gly Leu Met Gly Gly Gly Leu  
 25 65 70 75 80  
 Gly Gly Gly Leu Gly Asn Gly Leu Gly Gly Ser Gly Gly Leu Gly Glu  
 85 90 95  
 Gly Leu Ser Asn Ala Leu Asn Asp Met Leu Gly Gly Ser Leu Asn Thr  
 100 105 110  
 Leu Gly Ser Lys Gly Gly Asn Asn Thr Thr Ser Thr Thr Asn Ser Pro  
 30 115 120 125  
 Leu Asp Gln Ala Leu Gly Ile Asn Ser Thr Ser Gln Asn Asp Asp Ser  
 130 135 140  
 Thr Ser Gly Thr Asp Ser Thr Ser Asp Ser Ser Asp Pro Met Gln Gln  
 35 145 150 155 160  
 Leu Leu Lys Met Phe Ser Glu Ile Met Gln Ser Leu Phe Gly Asp Gly  
 165 170 175

- 8 -

	Gln	Asp	Gly	Thr	Gln	Gly	Ser	Ser	Ser	Gly	Gly	Lys	Gln	Pro	Thr	Glu	
				180					185					190			
	Gly	Glu	Gln	Asn	Ala	Tyr	Lys	Lys	Gly	Val	Thr	Asp	Ala	Leu	Ser	Gly	
			195					200					205				
5	Leu	Met	Gly	Asn	Gly	Leu	Ser	Gln	Leu	Leu	Gly	Asn	Gly	Gly	Leu	Gly	
		210					215					220					
	Gly	Gly	Gln	Gly	Gly	Asn	Ala	Gly	Thr	Gly	Leu	Asp	Gly	Ser	Ser	Leu	
		225				230					235					240	
10	Gly	Gly	Lys	Gly	Leu	Gln	Asn	Leu	Ser	Gly	Pro	Val	Asp	Tyr	Gln	Gln	
					245					250					255		
	Leu	Gly	Asn	Ala	Val	Gly	Thr	Gly	Ile	Gly	Met	Lys	Ala	Gly	Ile	Gln	
				260					265					270			
	Ala	Leu	Asn	Asp	Ile	Gly	Thr	His	Arg	His	Ser	Ser	Thr	Arg	Ser	Phe	
			275					280					285				
15	Val	Asn	Lys	Gly	Asp	Arg	Ala	Met	Ala	Lys	Glu	Ile	Gly	Gln	Phe	Met	
		290					295					300					
	Asp	Gln	Tyr	Pro	Glu	Val	Phe	Gly	Lys	Pro	Gln	Tyr	Gln	Lys	Gly	Pro	
		305				310					315					320	
20	Gly	Gln	Glu	Val	Lys	Thr	Asp	Asp	Lys	Ser	Trp	Ala	Lys	Ala	Leu	Ser	
					325					330					335		
	Lys	Pro	Asp	Asp	Asp	Gly	Met	Thr	Pro	Ala	Ser	Met	Glu	Gln	Phe	Asn	
				340					345					350			
	Lys	Ala	Lys	Gly	Met	Ile	Lys	Arg	Pro	Met	Ala	Gly	Asp	Thr	Gly	Asn	
			355					360					365				
25	Gly	Asn	Leu	Gln	Ala	Arg	Gly	Ala	Gly	Gly	Ser	Ser	Leu	Gly	Ile	Asp	
		370					375					380					
	Ala	Met	Met	Ala	Gly	Asp	Ala	Ile	Asn	Asn	Met	Ala	Leu	Gly	Lys	Leu	
		385				390				395						400	
	Gly	Ala	Ala														

30

This hypersensitive response elicitor polypeptide or protein has a molecular weight of about 39 kDa, has a pI of approximately 4.3, and is heat stable at 100°C for at least 10 minutes. This hypersensitive response elicitor polypeptide or protein has substantially no cysteine. The hypersensitive response elicitor polypeptide or protein derived from

35 *Erwinia amylovora* is more fully described in Wei, Z.-M., R. J. Laby, C. H. Zumoff, D. W. Bauer, S.-Y. He, A. Collmer, and S. V. Beer, "Harpin, Elicitor of the Hypersensitive Response Produced by the Plant Pathogen *Erwinia amylovora*,"

Science 257:85-88 (1992), which is hereby incorporated by reference. The DNA molecule encoding this polypeptide or protein has a nucleotide sequence corresponding to SEQ. ID. No. 4 as follows:

5	AAGCTTCGGC ATGGCACGTT TGACCGTTGG GTCGGCAGGG TACGTTTGAA TTATTCATAA	60
	GAGGAATACG TTATGAGTCT GAATACAAGT GGGCTGGGAG CGTCAACGAT GCAAATTTCT	120
	ATCGGCGGTG CGGGCGGAAA TAACGGGTTG CTGGGTACCA GTCGCCAGAA TGCTGGGTTG	180
	GGTGGCAATT CTGCACTGGG GCTGGGCGGC GGTAATCAAA ATGATACCGT CAATCAGCTG	240
	GCTGGCTTAC TCACCGGCAT GATGATGATG ATGAGCATGA TGGGCGGTGG TGGGCTGATG	300
10	GGCGGTGGCT TAGGCGGTGG CTTAGGTAAT GGCTTGGGTG GCTCAGGTGG CCTGGGCGAA	360
	GGACTGTCGA ACGCGCTGAA CGATATGTTA GGCGGTTTCG TGAACACGCT GGGCTCGAAA	420
	GGCGGCAACA ATACCACTTC AACAACAAAT TCCCCGCTGG ACCAGGCGCT GGGTATTAAC	480
	TCAACGTCCC AAAACGACGA TTCCACCTCC GGCACAGATT CCACCTCAGA CTCCAGCGAC	540
	CCGATGCAGC AGCTGCTGAA GATGTTGAGC GAGATAATGC AAAGCCTGTT TGGTGATGGG	600
15	CAAGATGGCA CCCAGGGCAG TTCCTCTGGG GGCAAGCAGC CGACCGAAGG CGAGCAGAAC	660
	GCCTATAAAA AAGGAGTCAC TGATGCGCTG TCGGCCTGA TGGGTAATGG TCTGAGCCAG	720
	CTCCTTGGCA ACGGGGGACT GGGAGGTGGT CAGGGCGGTA ATGCTGGCAC GGGTCTTGAC	780
	GGTTCGTCGC TGGGCGGCAA AGGGCTGCAA AACCTGAGCG GGCCGGTGGA CTACCAGCAG	840
	TTAGGTAACG CCGTGGGTAC CGGTATCGGT ATGAAAGCGG GCATTCAGGC GCTGAATGAT	900
20	ATCGGTACGC ACAGGCACAG TTCAACCCGT TCTTTCTGTC ATAAAGGCGA TCGGGCGATG	960
	GCGAAGGAAA TCGGTCAGTT CATGGACCAG TATCCTGAGG TGTTTGGCAA GCCGCAGTAC	1020
	CAGAAAGGCC CGGGTCAGGA GGTGAAAACC GATGACAAAT CATGGGCAA AGCACTGAGC	1080
	AAGCCAGATG ACGACGGAAT GACACCAGCC AGTATGGAGC AGTTCAACAA AGCCAAGGGC	1140
	ATGATCAAAA GGCCCATGGC GGGTGATACC GGCAACGGCA ACCTGCAGGC ACGCGGTGCC	1200
25	GGTGGTTCTT CGCTGGGTAT TGATGCCATG ATGGCCGGTG ATGCCATTAA CAATATGGCA	1260
	CTTGGCAAGC TGGGCGCGGC TTAAGCTT	1288

Another potentially suitable hypersensitive response elicitor from *Erwinia amylovora* is disclosed in U.S. Patent Application Serial No. 09/120,927, which is hereby incorporated by reference. The protein is encoded by a DNA molecule having a nucleic acid sequence of SEQ. ID. No. 5 as follows:

- 10 -

ATGTCAATTC TTACGCTTAA CAACAATACC TCGTCCTCGC CGGGTCTGTT CCAGTCCGGG 60  
 5 GGGGACAACG GGCTTGGTGG TCATAATGCA AATTCTGCGT TGGGGCAACA ACCCATCGAT 120  
 CGGCAAACCA TTGAGCAAAT GGCTCAATTA TTGGCGGAAC TGTTAAAGTC ACTGCTATCG 180  
 CCACAATCAG GTAATGCGGC AACCGGAGCC GGTGGCAATG ACCAGACTAC AGGAGTTGGT 240  
 10 AACGCTGGCG GCCTGAACGG ACGAAAAGGC ACAGCAGGAA CCACTCCGCA GTCTGACAGT 300  
 CAGAACATGC TGAGTGAGAT GGGCAACAAC GGGCTGGATC AGGCCATCAC GCCCGATGGC 360  
 15 CAGGGCGGGC GGCAGATCGG CGATAATCCT TTAAGTAAAG CCATGCTGAA GCTTATTGCA 420  
 CGCATGATGG ACGGCCAAAG CGATCAGTTT GGCCAACCTG GTACGGGCAA CAACAGTGCC 480  
 TCTTCCGGTA CTTCTTCATC TGGCGGTTCC CCTTTTAACG ATCTATCAGG GGGGAAGGCC 540  
 20 CCTTCCGGCA ACTCCCTTC CGGCAACTAC TCTCCCGTCA GTACCTTCTC ACCCCCATCC 600  
 ACGCCAACGT CCCCTACCTC ACCGCTTGAT TTCCCTTCTT CTCCACCAA AGCAGCCGGG 660  
 25 GGCAGCACGC CGGTAACCGA TCATCCTGAC CCTGTGGTA GCGCGGGCAT CGGGGCCGGA 720  
 AATTCGGTGG CCTTCACCAG CGCCGGCGCT AATCAGACGG TGCTGCATGA CACCATTACC 780  
 GTGAAAGCGG GTCAGGTGTT TGATGGCAAA GGACAAACCT TCACCGCCGG TTCAGAATTA 840  
 30 GGCGATGGCG GCCAGTCTGA AAACCAGAAA CCGCTGTTTA TACTGGAAGA CGGTGCCAGC 900  
 CTGAAAAACG TCACCATGGG CGACGACGGG GCGGATGGTA TTCATCTTTA CGGTGATGCC 960  
 35 AAAATAGACA ATCTGCACGT CACCAACGTG GGTGAGGACG CGATTACCGT TAAGCCAAAC 1020  
 AGCGCGGGCA AAAAATCCCA CGTTGAAATC ACTAACAGTT CCTTCGAGCA CGCCTCTGAC 1080  
 AAGATCCTGC AGCTGAATGC CGATACTAAC CTGAGCGTTG ACAACGTGAA GGCCAAAGAC 1140  
 40 TTTGGTACTT TTGTACGCAC TAACGGCGGT CAACAGGGTA ACTGGGATCT GAATCTGAGC 1200  
 CATATCAGCG CAGAAGACGG TAAGTTCTCG TTCGTTAAAA GCGATAGCGA GGGGCTAAAC 1260  
 45 GTCAATACCA GTGATATCTC ACTGGGTGAT GTTGAAAACC ACTACAAAGT GCCGATGTCC 1320  
 GCCAACCTGA AGGTGGCTGA ATGA 1344

See GenBank Accession No. U94513. The isolated DNA molecule of the present  
 50 invention encodes a hypersensitive response elicitor protein or polypeptide having an  
 amino acid sequence of SEQ. ID. No. 6 as follows:

Met Ser Ile Leu Thr Leu Asn Asn Asn Thr Ser Ser Ser Pro Gly Leu  
 1 5 10 15  
 55 Phe Gln Ser Gly Gly Asp Asn Gly Leu Gly Gly His Asn Ala Asn Ser  
 20 25 30  
 60 Ala Leu Gly Gln Gln Pro Ile Asp Arg Gln Thr Ile Glu Gln Met Ala  
 35 40 45

- 11 -

	Gln	Leu	Leu	Ala	Glu	Leu	Leu	Lys	Ser	Leu	Leu	Ser	Pro	Gln	Ser	Gly	
	50						55					60					
5	Asn	Ala	Ala	Thr	Gly	Ala	Gly	Gly	Asn	Asp	Gln	Thr	Thr	Gly	Val	Gly	
	65					70				75						80	
	Asn	Ala	Gly	Gly	Leu	Asn	Gly	Arg	Lys	Gly	Thr	Ala	Gly	Thr	Thr	Pro	
					85					90					95		
10	Gln	Ser	Asp	Ser	Gln	Asn	Met	Leu	Ser	Glu	Met	Gly	Asn	Asn	Gly	Leu	
				100					105					110			
	Asp	Gln	Ala	Ile	Thr	Pro	Asp	Gly	Gln	Gly	Gly	Gly	Gln	Ile	Gly	Asp	
15			115					120					125				
	Asn	Pro	Leu	Leu	Lys	Ala	Met	Leu	Lys	Leu	Ile	Ala	Arg	Met	Met	Asp	
		130					135					140					
20	Gly	Gln	Ser	Asp	Gln	Phe	Gly	Gln	Pro	Gly	Thr	Gly	Asn	Asn	Ser	Ala	
	145					150					155					160	
	Ser	Ser	Gly	Thr	Ser	Ser	Ser	Gly	Gly	Ser	Pro	Phe	Asn	Asp	Leu	Ser	
					165					170					175		
25	Gly	Gly	Lys	Ala	Pro	Ser	Gly	Asn	Ser	Pro	Ser	Gly	Asn	Tyr	Ser	Pro	
				180					185					190			
	Val	Ser	Thr	Phe	Ser	Pro	Pro	Ser	Thr	Pro	Thr	Ser	Pro	Thr	Ser	Pro	
30			195					200					205				
	Leu	Asp	Phe	Pro	Ser	Ser	Pro	Thr	Lys	Ala	Ala	Gly	Gly	Ser	Thr	Pro	
		210					215					220					
35	Val	Thr	Asp	His	Pro	Asp	Pro	Val	Gly	Ser	Ala	Gly	Ile	Gly	Ala	Gly	
	225					230					235					240	
	Asn	Ser	Val	Ala	Phe	Thr	Ser	Ala	Gly	Ala	Asn	Gln	Thr	Val	Leu	His	
				245						250					255		
40	Asp	Thr	Ile	Thr	Val	Lys	Ala	Gly	Gln	Val	Phe	Asp	Gly	Lys	Gly	Gln	
				260					265					270			
	Thr	Phe	Thr	Ala	Gly	Ser	Glu	Leu	Gly	Asp	Gly	Gly	Gln	Ser	Glu	Asn	
45			275					280					285				
	Gln	Lys	Pro	Leu	Phe	Ile	Leu	Glu	Asp	Gly	Ala	Ser	Leu	Lys	Asn	Val	
		290					295					300					
50	Thr	Met	Gly	Asp	Asp	Gly	Ala	Asp	Gly	Ile	His	Leu	Tyr	Gly	Asp	Ala	
	305				310						315					320	
	Lys	Ile	Asp	Asn	Leu	His	Val	Thr	Asn	Val	Gly	Glu	Asp	Ala	Ile	Thr	
				325						330					335		
55	Val	Lys	Pro	Asn	Ser	Ala	Gly	Lys	Lys	Ser	His	Val	Glu	Ile	Thr	Asn	
				340					345					350			

- 12 -

Ser Ser Phe Glu His Ala Ser Asp Lys Ile Leu Gln Leu Asn Ala Asp  
 355 360 365  
 5 Thr Asn Leu Ser Val Asp Asn Val Lys Ala Lys Asp Phe Gly Thr Phe  
 370 375 380  
 Val Arg Thr Asn Gly Gly Gln Gln Gly Asn Trp Asp Leu Asn Leu Ser  
 385 390 395 400  
 10 His Ile Ser Ala Glu Asp Gly Lys Phe Ser Phe Val Lys Ser Asp Ser  
 405 410 415  
 Glu Gly Leu Asn Val Asn Thr Ser Asp Ile Ser Leu Gly Asp Val Glu  
 420 425 430  
 15 Asn His Tyr Lys Val Pro Met Ser Ala Asn Leu Lys Val Ala Glu  
 435 440 445

20 This protein or polypeptide is acidic, rich in glycine and serine, and lacks cysteine. It is also heat stable, protease sensitive, and suppressed by inhibitors of plant metabolism. The protein or polypeptide of the present invention has a predicted molecular size of ca. 4.5 kDa.

Another potentially suitable hypersensitive response elicitor from  
 25 *Erwinia amylovora* is disclosed in U.S. Patent Application Serial No. 09/120,663, which is hereby incorporated by reference. The protein is encoded by a DNA molecule having a nucleic acid sequence of SEQ. ID. No. 7 as follows:

30 ATGGAATTAA AATCACTGGG AACTGAACAC AAGGCGGCAG TACACACAGC GGCGCACAAAC 60  
 CCTGTGGGGC ATGGTGTTCG CTTACAGCAG GGCAGCAGCA GCAGCAGCCC GCAAATGCC 120  
 GCTGCATCAT TGGCGGCAGA AGGCAAAAAT CGTGGGAAAA TGCCGAGAAT TCACCAGCCA 180  
 35 TCTACTGCGG CTGATGGTAT CAGCGCTGCT CACCAGCAAA AGAAATCCTT CAGTCTCAGG 240  
 GGCTGTTTGG GGACGAAAAA ATTTTCCAGA TCGGCACCGC AGGGCCAGCC AGGTACCACC 300  
 CACAGCAAAG GGGCAACATT GCGCGATCTG CTGGCGCGGG ACGACGGCGA AACGCAGCAT 360  
 40 GAGGCGGCCG CGCCAGATGC GGCGCGTTTG ACCCGTTCGG GCGGCGTCAA ACGCCGCAAT 420  
 ATGGACGACA TGGCCGGGCG GCCAATGGTG AAAGGTGGCA GCGGCGAAGA TAAGGTACCA 480  
 45 ACGCAGCAAA AACGGCATCA GCTGAACAAT TTTGGCCAGA TGCGCCAAAC GATGTTGAGC 540  
 AAAATGGCTC ACCCGGCTTC AGCCAACGCC GGCATCGCC TGCAGCATTC ACCGCCGCAC 600  
 ATCCCGGGTA GCCACCACGA AATCAAGGAA GAACCGGTTG GCTCCACCAG CAAGGCAACA 660  
 50 ACGGCCCCAG CAGACAGAGT GGAAATCGCT CAGGAAGATG ACGACAGCGA ATTCCAGCAA 720  
 CTGCATCAAC AGCGGCTGGC GCGCGAACGG GAAAATCCAC CGCAGCCGCC CAAACTCGGC 780  
 55 GTTGCCACAC CGATTAGCGC CAGGTTTCAG CCCAACTGA CTGCGGTTGC GGAAAGCGTC 840

	CTTGAGGGGA	CAGATACCAC	GCAGTCACCC	CTTAAGCCGC	AATCAATGCT	GAAAGGAAGT	900
5	GGAGCCGGGG	TAACGCCGCT	GGCGGTAACG	CTGGATAAAG	GCAAGTTGCA	GCTGGCACCG	960
	GATAATCCAC	CCGCGCTCAA	TACGTTGTTG	AAGCAGACAT	TGGGTAAAGA	CACCCAGCAC	1020
	TATCTGGCGC	ACCATGCCAG	CAGCGACGGT	AGCCAGCATC	TGCTGCTGGA	CAACAAAGGC	1080
10	CACCTGTTTG	ATATCAAAAG	CACCGCCACC	AGCTATAGCG	TGCTGCACAA	CAGCCACCCC	1140
	GGTGAGATAA	AGGGCAAGCT	GGCGCAGGCG	GGTACTGGCT	CCGTCAGCGT	AGACGGTAAA	1200
15	AGCGGCAAGA	TCTCGCTGGG	GAGCGGTACG	CAAAGTCACA	ACAAAACAAT	GCTAAGCCAA	1260
	CCGGGGGAAG	CGCACCGTTC	CTTATTAACC	GGCATTGGC	AGCATCCTGC	TGGCGCAGCG	1320
	CGGCCGCAGG	GCGAGTCAAT	CCGCCTGCAT	GACGACAAAA	TTCATATCCT	GCATCCGGAG	1380
20	CTGGGCGTAT	GGCAATCTGC	GGATAAAGAT	ACCCACAGCC	AGCTGTCTCG	CCAGGCAGAC	1440
	GGTAAGCTCT	ATGCGCTGAA	AGACAACCGT	ACCTGCAAA	ACCTCTCCGA	TAATAAATCC	1500
25	TCAGAAAAGC	TGTCGATAA	AATCAAATCG	TATTCCGTTG	ATCAGCGGGG	GCAGGTGGCG	1560
	ATCCTGACGG	ATACTCCCGG	CCGCCATAAG	ATGAGTATTA	TGCCCTCGCT	GGATGCTTCC	1620
	CCGGAGAGCC	ATATTTCCCT	CAGCCTGCAT	TTTGCCGATG	CCCACCAGGG	GTTATTGCAC	1680
30	GGGAAGTCGG	AGCTTGAGGC	ACAATCTGTC	GCGATCAGCC	ATGGGCGACT	GGTTGTGGCC	1740
	GATAGCGAAG	GCAAGCTGTT	TAGCGCCGCC	ATTCGGAAGC	AAGGGGATGG	AAACGAAGTG	1800
35	AAAATGAAAG	CCATGCCTCA	GCATGCGCTC	GATGAACATT	TTGGTCATGA	CCACCAGATT	1860
	TCTGGATTTT	TCCATGACGA	CCACGGCCAG	CTTAATGCGC	TGGTGAAAAA	TAACCTCAGG	1920
	CAGCAGCATG	CCTGCCCCTT	GGGTAACGAT	CATCAGTTTC	ACCCCGGCTG	GAACCTGACT	1980
40	GATGCGCTGG	TTATCGACAA	TCAGCTGGGG	CTGCATCATA	CCAATCCTGA	ACCGCATGAG	2040
	ATTCTTGATA	TGGGGCATTT	AGGCAGCCTG	GCGTTACAGG	AGGGCAAGCT	TCACTATTTT	2100
45	GACCAGCTGA	CCAAAGGGTG	GA CTGGCGCG	GAGTCAGATT	GTAAGCAGCT	GAAAAAAGGC	2160
	CTGGATGGAG	CAGCTTATCT	ACTGAAAGAC	GGTGAAGTGA	AACGCCTGAA	TATTAATCAG	2220
	AGCACCTCCT	CTATCAAGCA	CGGAACGGAA	AACGTTTTTT	CGCTGCCGCA	TGTGCGCAAT	2280
50	AAACCGGAGC	CGGGAGATGC	CCTGCAAGGG	CTGAATAAAG	ACGATAAGGC	CCAGGCCATG	2340
	GCGGTGATTG	GGGTAAATAA	ATACCTGGCG	CTGACGGAAA	AAGGGGACAT	TCGCTCCTTC	2400
55	CAGATAAAAC	CCGGCACCCA	GCAGTTGGAG	CGGCCGGCAC	AAACTCTCAG	CCGCGAAGGT	2460
	ATCAGCGGCG	AACTGAAAGA	CATTCTATGTC	GACCACAAGC	AGAACCTGTA	TGCCTTGACC	2520
	CACGAGGGAG	AGGTGTTTCA	TCAGCCGCGT	GAAGCCTGGC	AGAATGGTGC	CGAAAGCAGC	2580
60	AGCTGGCACA	AACTGGCGTT	GCCACAGAGT	GAAAGTAAGC	TAAAAAGTCT	GGACATGAGC	2640
	CATGAGCACA	AACCGATTGC	CACCTTTGAA	GACGGTAGCC	AGCATCAGCT	GAAGGCTGGC	2700
65	GGCTGGCACG	CCTATGCGGC	ACCTGAACGC	GGGCCGCTGG	CGGTGGGTAC	CAGCGGTTCA	2760

- 14 -

	CAAACCGTCT	TTAACCGACT	AATGCAGGGG	GTGAAAGGCA	AGGTGATCCC	AGGCAGCGGG	2820
	TTGACGGTTA	AGCTCTCGGC	TCAGACGGGG	GGAATGACCG	GCGCCGAAGG	GCGCAAGGTC	2880
5	AGCAGTAAAT	TTTCCGAAAG	GATCCGCGCC	TATGCGTTCA	ACCCAACAAT	GTCCACGCCG	2940
	CGACCGATTA	AAAATGCTGC	TTATGCCACA	CAGCACGGCT	GGCAGGGGCG	TGAGGGGTTG	3000
10	AAGCCGTTGT	ACGAGATGCA	GGGAGCGCTG	ATTAAACAAC	TGGATGCGCA	TAACGTTCTG	3060
	CATAACGCGC	CACAGCCAGA	TTGTCAGAGC	AAACTGGAAA	CTCTGGATTT	AGGCGAACAT	3120
	GGCGCAGAAT	TGCTTAACGA	CATGAAGCGC	TTCCGCGACG	AACTGGAGCA	GAGTGCAACC	3180
15	CGTTCGGTGA	CCGTTTTAGG	TCAACATCAG	GGAGTGCTAA	AAAGCAACGG	TGAAATCAAT	3240
	AGCGAATTTA	AGCCATCGCC	CGGCAAGGCG	TTGGTCCAGA	GCTTTAACGT	CAATCGCTCT	3300
20	GGTCAGGATC	TAAGCAAGTC	ACTGCAACAG	GCAGTACATG	CCACGCCGCC	ATCCGCAGAG	3360
	AGTAAACTGC	AATCCATGCT	GGGGCACTTT	GTCAGTGCCG	GGGTGGATAT	GAGTCATCAG	3420
	AAGGGCGAGA	TCCCGCTGGG	CCGCCAGCGC	GATCCGAATG	ATAAAACCGC	ACTGACCAAA	3480
25	TCGCGTTTAA	TTTLAGATAC	CGTGACCATC	GGTGAAGTGC	ATGAAGTGGC	CGATAAGGCG	3540
	AAACTGGTAT	CTGACCATAA	ACCCGATGCC	GATCAGATAA	AACAGCTGCG	CCAGCAGTTC	3600
30	GATACGCTGC	GTGAAAAGCG	GTATGAGAGC	AATCCGGTGA	AGCATTACAC	CGATATGGGC	3660
	TTCACCCATA	ATAAGGCGCT	GGAAGCAAAC	TATGATGCGG	TCAAAGCCTT	TATCAATGCC	3720
	TTTAAGAAAG	AGCACCACGG	CGTCAATCTG	ACCACGCGTA	CCGTACTGGA	ATCACAGGGC	3780
35	AGTGCGGAGC	TGGCGAAGAA	GCTCAAGAAT	ACGCTGTTGT	CCCTGGACAG	TGGTGAAAGT	3840
	ATGAGCTTCA	GCCGGTCATA	TGGCGGGGGC	GTCAGCACTG	TCTTTGTGCC	TACCCCTAGC	3900
40	AAGAAGGTGC	CAGTTCCGGT	GATCCCCGGA	GCCGGCATCA	CGCTGGATCG	CGCCTATAAC	3960
	CTGAGCTTCA	GTCGTACCAG	CGGCGGATTG	AACGTCAGTT	TTGGCCGCGA	CGGCGGGGTG	4020
	AGTGGTAACA	TCATGGTCGC	TACCGGCCAT	GATGTGATGC	CCTATATGAC	CGGTAAGAAA	4080
45	ACCAGTGCAG	GTAACGCCAG	TGACTGGTTG	AGCGCAAAAC	ATAAAATCAG	CCCGGACTTG	4140
	CGTATCGGCG	CTGCTGTGAG	TGGCACCCCTG	CAAGGAACGC	TACAAAACAG	CCTGAAGTTT	4200
50	AAGCTGACAG	AGGATGAGCT	GCCTGGCTTT	ATCCATGGCT	TGACGCATGG	CACGTTGACC	4260
	CCGGCAGAAC	TGTTGCAAAA	GGGGATCGAA	CATCAGATGA	AGCAGGGCAG	CAAACGTACG	4320
	TTTAGCGTCG	ATACCTCGGC	AAATCTGGAT	CTGCGTGCCG	GTATCAATCT	GAACGAAGAC	4380
55	GGCAGTAAAC	CAAATGGTGT	CACTGCCCGT	GTTTCTGCCG	GGCTAAGTGC	ATCGGCAAAC	4440
	CTGGCCGCCG	GCTCGCGTGA	ACGCAGCACC	ACCTCTGGCC	AGTTTGGCAG	CACGACTTCG	4500
60	GCCAGCAATA	ACCGCCCAAC	CTTCCTCAAC	GGGGTCGGCG	CGGGTGCTAA	CCTGACGGCT	4560
	GCTTTAGGGG	TTGCCCATTC	ATCTACGCAT	GAAGGGAAAC	CGGTGCGGAT	CTTCCCGGCA	4620
	TTTACCTCGA	CCAATGTTTC	GGCAGCGCTG	GCGCTGGATA	ACCGTACCTC	ACAGAGTATC	4680
65	AGCCTGGAAT	TGAAGCGCGC	GGAGCCGGTG	ACCAGCAACG	ATATCAGCGA	GTTGACCTCC	4740



- 15 -

ACGCTGGGAA AACACTTTAA GGATAGCGCC ACAACGAAGA TGCTTGCCGC TCTCAAAGAG 4800  
 TTAGATGACG CTAAGCCCGC TGAACAACTG CATATTTTAC AGCAGCATTT CAGTGCAAAA 4860  
 5 GATGTCGTCG GTGATGAACG CTACGAGGCG GTGCGCAACC TGAAAAAACT GGTGATACGT 4920  
 CAACAGGCTG CGGACAGCCA CAGCATGGAA TTAGGATCTG CCAGTCACAG CACGACCTAC 4980  
 10 AATAATCTGT CGAGAATAAA TAATGACGGC ATTGTCGAGC TGCTACACAA ACATTTCGAT 5040  
 GCGGCATTAC CAGCAAGCAG TGCCAAACGT CTTGGTGAAA TGATGAATAA CGATCCGGCA 5100  
 CTGAAAGATA TTATTAAGCA GCTGCAAAGT ACGCCGTTCA GCAGCGCCAG CGTGTGATG 5160  
 15 GAGCTGAAAG ATGGTCTGCG TGAGCAGACG GAAAAAGCAA TACTGGACGG TAAGGTCGGT 5220  
 CGTGAAGAAG TGGGAGTACT TTTCCAGGAT CGTAACAACT TGCGTGTTAA ATCGGTCAGC 5280  
 20 GTCAGTCAGT CCGTCAGCAA AAGCGAAGGC TTCAATACCC CAGCGCTGTT ACTGGGGACG 5340  
 AGCAACAGCG CTGCTATGAG CATGGAGCGC AACATCGGAA CCATTAATTT TAAATACGGC 5400  
 CAGGATCAGA ACACCCACG GCGATTTACC CTGGAGGGTG GAATAGCTCA GGCTAATCCG 5460  
 25 CAGGTCGCAT CTGCGCTTAC TGATTGTAAG AAGGAAGGGC TGGAAATGAA GAGCTAA 5517

This DNA molecule is known as the dspE gene for *Erwinia amylovora*. This isolated  
 30 DNA molecule of the present invention encodes a protein or polypeptide which elicits  
 a plant pathogen's hypersensitive response having an amino acid sequence of SEQ.  
 ID. No. 8 as follows:

35 Met Glu Leu Lys Ser Leu Gly Thr Glu His Lys Ala Ala Val His Thr  
 1 5 10 15  
 Ala Ala His Asn Pro Val Gly His Gly Val Ala Leu Gln Gln Gly Ser  
 20 25 30  
 40 Ser Ser Ser Ser Pro Gln Asn Ala Ala Ala Ser Leu Ala Ala Glu Gly  
 35 40 45  
 Lys Asn Arg Gly Lys Met Pro Arg Ile His Gln Pro Ser Thr Ala Ala  
 50 55 60  
 45 Asp Gly Ile Ser Ala Ala His Gln Gln Lys Lys Ser Phe Ser Leu Arg  
 65 70 75 80  
 Gly Cys Leu Gly Thr Lys Lys Phe Ser Arg Ser Ala Pro Gln Gly Gln  
 85 90 95  
 50 Pro Gly Thr Thr His Ser Lys Gly Ala Thr Leu Arg Asp Leu Leu Ala  
 100 105 110  
 55 Arg Asp Asp Gly Glu Thr Gln His Glu Ala Ala Ala Pro Asp Ala Ala  
 115 120 125  
 Arg Leu Thr Arg Ser Gly Gly Val Lys Arg Arg Asn Met Asp Asp Met  
 130 135 140  
 60

- 16 -

	Ala Gly Arg Pro Met Val Lys Gly Gly Ser Gly Glu Asp Lys Val Pro	145	150	155	160
5	Thr Gln Gln Lys Arg His Gln Leu Asn Asn Phe Gly Gln Met Arg Gln	165	170	175	
	Thr Met Leu Ser Lys Met Ala His Pro Ala Ser Ala Asn Ala Gly Asp	180	185	190	
10	Arg Leu Gln His Ser Pro Pro His Ile Pro Gly Ser His His Glu Ile	195	200	205	
	Lys Glu Glu Pro Val Gly Ser Thr Ser Lys Ala Thr Thr Ala His Ala	210	215	220	
15	Asp Arg Val Glu Ile Ala Gln Glu Asp Asp Asp Ser Glu Phe Gln Gln	225	230	235	240
	Leu His Gln Gln Arg Leu Ala Arg Glu Arg Glu Asn Pro Pro Gln Pro	245	250	255	
20	Pro Lys Leu Gly Val Ala Thr Pro Ile Ser Ala Arg Phe Gln Pro Lys	260	265	270	
	Leu Thr Ala Val Ala Glu Ser Val Leu Glu Gly Thr Asp Thr Thr Gln	275	280	285	
25	Ser Pro Leu Lys Pro Gln Ser Met Leu Lys Gly Ser Gly Ala Gly Val	290	295	300	
30	Thr Pro Leu Ala Val Thr Leu Asp Lys Gly Lys Leu Gln Leu Ala Pro	305	310	315	320
	Asp Asn Pro Pro Ala Leu Asn Thr Leu Leu Lys Gln Thr Leu Gly Lys	325	330	335	
35	Asp Thr Gln His Tyr Leu Ala His His Ala Ser Ser Asp Gly Ser Gln	340	345	350	
40	His Leu Leu Leu Asp Asn Lys Gly His Leu Phe Asp Ile Lys Ser Thr	355	360	365	
	Ala Thr Ser Tyr Ser Val Leu His Asn Ser His Pro Gly Glu Ile Lys	370	375	380	
45	Gly Lys Leu Ala Gln Ala Gly Thr Gly Ser Val Ser Val Asp Gly Lys	385	390	395	400
	Ser Gly Lys Ile Ser Leu Gly Ser Gly Thr Gln Ser His Asn Lys Thr	405	410	415	
50	Met Leu Ser Gln Pro Gly Glu Ala His Arg Ser Leu Leu Thr Gly Ile	420	425	430	
	Trp Gln His Pro Ala Gly Ala Ala Arg Pro Gln Gly Glu Ser Ile Arg	435	440	445	
55	Leu His Asp Asp Lys Ile His Ile Leu His Pro Glu Leu Gly Val Trp	450	455	460	
60	Gln Ser Ala Asp Lys Asp Thr His Ser Gln Leu Ser Arg Gln Ala Asp	465	470	475	480
	Gly Lys Leu Tyr Ala Leu Lys Asp Asn Arg Thr Leu Gln Asn Leu Ser	485	490	495	
65					

- 17 -

	Asp Asn Lys Ser Ser Glu Lys Leu Val Asp Lys Ile Lys Ser Tyr Ser	500	505	510
5	Val Asp Gln Arg Gly Gln Val Ala Ile Leu Thr Asp Thr Pro Gly Arg	515	520	525
	His Lys Met Ser Ile Met Pro Ser Leu Asp Ala Ser Pro Glu Ser His	530	535	540
10	Ile Ser Leu Ser Leu His Phe Ala Asp Ala His Gln Gly Leu Leu His	545	550	555
	Gly Lys Ser Glu Leu Glu Ala Gln Ser Val Ala Ile Ser His Gly Arg	565	570	575
15	Leu Val Val Ala Asp Ser Glu Gly Lys Leu Phe Ser Ala Ala Ile Pro	580	585	590
	Lys Gln Gly Asp Gly Asn Glu Leu Lys Met Lys Ala Met Pro Gln His	595	600	605
20	Ala Leu Asp Glu His Phe Gly His Asp His Gln Ile Ser Gly Phe Phe	610	615	620
	His Asp Asp His Gly Gln Leu Asn Ala Leu Val Lys Asn Asn Phe Arg	625	630	635
25	Gln Gln His Ala Cys Pro Leu Gly Asn Asp His Gln Phe His Pro Gly	645	650	655
30	Trp Asn Leu Thr Asp Ala Leu Val Ile Asp Asn Gln Leu Gly Leu His	660	665	670
	His Thr Asn Pro Glu Pro His Glu Ile Leu Asp Met Gly His Leu Gly	675	680	685
35	Ser Leu Ala Leu Gln Glu Gly Lys Leu His Tyr Phe Asp Gln Leu Thr	690	695	700
	Lys Gly Trp Thr Gly Ala Glu Ser Asp Cys Lys Gln Leu Lys Lys Gly	705	710	715
40	Leu Asp Gly Ala Ala Tyr Leu Leu Lys Asp Gly Glu Val Lys Arg Leu	725	730	735
45	Asn Ile Asn Gln Ser Thr Ser Ser Ile Lys His Gly Thr Glu Asn Val	740	745	750
	Phe Ser Leu Pro His Val Arg Asn Lys Pro Glu Pro Gly Asp Ala Leu	755	760	765
50	Gln Gly Leu Asn Lys Asp Asp Lys Ala Gln Ala Met Ala Val Ile Gly	770	775	780
	Val Asn Lys Tyr Leu Ala Leu Thr Glu Lys Gly Asp Ile Arg Ser Phe	785	790	795
55	Gln Ile Lys Pro Gly Thr Gln Gln Leu Glu Arg Pro Ala Gln Thr Leu	805	810	815
60	Ser Arg Glu Gly Ile Ser Gly Glu Leu Lys Asp Ile His Val Asp His	820	825	830
	Lys Gln Asn Leu Tyr Ala Leu Thr His Glu Gly Glu Val Phe His Gln	835	840	845
65				

- 18 -

	Pro	Arg	Glu	Ala	Trp	Gln	Asn	Gly	Ala	Glu	Ser	Ser	Ser	Trp	His	Lys	
	850						855					860					
5	Leu	Ala	Leu	Pro	Gln	Ser	Glu	Ser	Lys	Leu	Lys	Ser	Leu	Asp	Met	Ser	
	865					870					875					880	
	His	Glu	His	Lys	Pro	Ile	Ala	Thr	Phe	Glu	Asp	Gly	Ser	Gln	His	Gln	
					885					890					895		
10	Leu	Lys	Ala	Gly	Gly	Trp	His	Ala	Tyr	Ala	Ala	Pro	Glu	Arg	Gly	Pro	
				900					905					910			
	Leu	Ala	Val	Gly	Thr	Ser	Gly	Ser	Gln	Thr	Val	Phe	Asn	Arg	Leu	Met	
			915					920					925				
15	Gln	Gly	Val	Lys	Gly	Lys	Val	Ile	Pro	Gly	Ser	Gly	Leu	Thr	Val	Lys	
	930						935					940					
	Leu	Ser	Ala	Gln	Thr	Gly	Gly	Met	Thr	Gly	Ala	Glu	Gly	Arg	Lys	Val	
20	945					950					955					960	
	Ser	Ser	Lys	Phe	Ser	Glu	Arg	Ile	Arg	Ala	Tyr	Ala	Phe	Asn	Pro	Thr	
					965					970					975		
25	Met	Ser	Thr	Pro	Arg	Pro	Ile	Lys	Asn	Ala	Ala	Tyr	Ala	Thr	Gln	His	
				980					985					990			
	Gly	Trp	Gln	Gly	Arg	Glu	Gly	Leu	Lys	Pro	Leu	Tyr	Glu	Met	Gln	Gly	
			995					1000					1005				
30	Ala	Leu	Ile	Lys	Gln	Leu	Asp	Ala	His	Asn	Val	Arg	His	Asn	Ala	Pro	
	1010						1015					1020					
	Gln	Pro	Asp	Leu	Gln	Ser	Lys	Leu	Glu	Thr	Leu	Asp	Leu	Gly	Glu	His	
35	1025					1030					1035					1040	
	Gly	Ala	Glu	Leu	Leu	Asn	Asp	Met	Lys	Arg	Phe	Arg	Asp	Glu	Leu	Glu	
					1045					1050					1055		
40	Gln	Ser	Ala	Thr	Arg	Ser	Val	Thr	Val	Leu	Gly	Gln	His	Gln	Gly	Val	
					1060				1065					1070			
	Leu	Lys	Ser	Asn	Gly	Glu	Ile	Asn	Ser	Glu	Phe	Lys	Pro	Ser	Pro	Gly	
			1075					1080					1085				
45	Lys	Ala	Leu	Val	Gln	Ser	Phe	Asn	Val	Asn	Arg	Ser	Gly	Gln	Asp	Leu	
	1090						1095					1100					
	Ser	Lys	Ser	Leu	Gln	Gln	Ala	Val	His	Ala	Thr	Pro	Pro	Ser	Ala	Glu	
50	1105					1110					1115					1120	
	Ser	Lys	Leu	Gln	Ser	Met	Leu	Gly	His	Phe	Val	Ser	Ala	Gly	Val	Asp	
					1125					1130					1135		
55	Met	Ser	His	Gln	Lys	Gly	Glu	Ile	Pro	Leu	Gly	Arg	Gln	Arg	Asp	Pro	
				1140					1145					1150			
	Asn	Asp	Lys	Thr	Ala	Leu	Thr	Lys	Ser	Arg	Leu	Ile	Leu	Asp	Thr	Val	
			1155					1160					1165				
60	Thr	Ile	Gly	Glu	Leu	His	Glu	Leu	Ala	Asp	Lys	Ala	Lys	Leu	Val	Ser	
							1175					1180					
	Asp	His	Lys	Pro	Asp	Ala	Asp	Gln	Ile	Lys	Gln	Leu	Arg	Gln	Gln	Phe	
65	1185					1190					1195					1200	

- 19 -

	Asp Thr Leu Arg Glu Lys Arg Tyr Glu Ser Asn Pro Val Lys His Tyr	
	1205	1210 1215
5	Thr Asp Met Gly Phe Thr His Asn Lys Ala Leu Glu Ala Asn Tyr Asp	
	1220	1225 1230
	Ala Val Lys Ala Phe Ile Asn Ala Phe Lys Lys Glu His His Gly Val	
	1235	1240 1245
10	Asn Leu Thr Thr Arg Thr Val Leu Glu Ser Gln Gly Ser Ala Glu Leu	
	1250	1255 1260
	Ala Lys Lys Leu Lys Asn Thr Leu Leu Ser Leu Asp Ser Gly Glu Ser	
15	1265	1270 1275 1280
	Met Ser Phe Ser Arg Ser Tyr Gly Gly Gly Val Ser Thr Val Phe Val	
	1285	1290 1295
20	Pro Thr Leu Ser Lys Lys Val Pro Val Pro Val Ile Pro Gly Ala Gly	
	1300	1305 1310
	Ile Thr Leu Asp Arg Ala Tyr Asn Leu Ser Phe Ser Arg Thr Ser Gly	
	1315	1320 1325
25	Gly Leu Asn Val Ser Phe Gly Arg Asp Gly Gly Val Ser Gly Asn Ile	
	1330	1335 1340
	Met Val Ala Thr Gly His Asp Val Met Pro Tyr Met Thr Gly Lys Lys	
30	1345	1350 1355 1360
	Thr Ser Ala Gly Asn Ala Ser Asp Trp Leu Ser Ala Lys His Lys Ile	
	1365	1370 1375
35	Ser Pro Asp Leu Arg Ile Gly Ala Ala Val Ser Gly Thr Leu Gln Gly	
	1380	1385 1390
	Thr Leu Gln Asn Ser Leu Lys Phe Lys Leu Thr Glu Asp Glu Leu Pro	
	1395	1400 1405
40	Gly Phe Ile His Gly Leu Thr His Gly Thr Leu Thr Pro Ala Glu Leu	
	1410	1415 1420
	Leu Gln Lys Gly Ile Glu His Gln Met Lys Gln Gly Ser Lys Leu Thr	
45	1425	1430 1435 1440
	Phe Ser Val Asp Thr Ser Ala Asn Leu Asp Leu Arg Ala Gly Ile Asn	
	1445	1450 1455
50	Leu Asn Glu Asp Gly Ser Lys Pro Asn Gly Val Thr Ala Arg Val Ser	
	1460	1465 1470
	Ala Gly Leu Ser Ala Ser Ala Asn Leu Ala Ala Gly Ser Arg Glu Arg	
	1475	1480 1485
55	Ser Thr Thr Ser Gly Gln Phe Gly Ser Thr Thr Ser Ala Ser Asn Asn	
	1490	1495 1500
	Arg Pro Thr Phe Leu Asn Gly Val Gly Ala Gly Ala Asn Leu Thr Ala	
60	1505	1510 1515 1520
	Ala Leu Gly Val Ala His Ser Ser Thr His Glu Gly Lys Pro Val Gly	
	1525	1530 1535

- 20 -

	Ile Phe Pro Ala Phe Thr Ser Thr Asn Val Ser Ala Ala Leu Ala Leu	1540	1545	1550
5	Asp Asn Arg Thr Ser Gln Ser Ile Ser Leu Glu Leu Lys Arg Ala Glu	1555	1560	1565
	Pro Val Thr Ser Asn Asp Ile Ser Glu Leu Thr Ser Thr Leu Gly Lys	1570	1575	1580
10	His Phe Lys Asp Ser Ala Thr Thr Lys Met Leu Ala Ala Leu Lys Glu	1585	1590	1595
	Leu Asp Asp Ala Lys Pro Ala Glu Gln Leu His Ile Leu Gln Gln His	1605	1610	1615
15	Phe Ser Ala Lys Asp Val Val Gly Asp Glu Arg Tyr Glu Ala Val Arg	1620	1625	1630
	Asn Leu Lys Lys Leu Val Ile Arg Gln Gln Ala Ala Asp Ser His Ser	1635	1640	1645
20	Met Glu Leu Gly Ser Ala Ser His Ser Thr Thr Tyr Asn Asn Leu Ser	1650	1655	1660
	Arg Ile Asn Asn Asp Gly Ile Val Glu Leu Leu His Lys His Phe Asp	1665	1670	1675
25	Ala Ala Leu Pro Ala Ser Ser Ala Lys Arg Leu Gly Glu Met Met Asn	1685	1690	1695
	Asn Asp Pro Ala Leu Lys Asp Ile Ile Lys Gln Leu Gln Ser Thr Pro	1700	1705	1710
30	Phe Ser Ser Ala Ser Val Ser Met Glu Leu Lys Asp Gly Leu Arg Glu	1715	1720	1725
	Gln Thr Glu Lys Ala Ile Leu Asp Gly Lys Val Gly Arg Glu Glu Val	1730	1735	1740
35	Gly Val Leu Phe Gln Asp Arg Asn Asn Leu Arg Val Lys Ser Val Ser	1745	1750	1755
	Val Ser Gln Ser Val Ser Lys Ser Glu Gly Phe Asn Thr Pro Ala Leu	1765	1770	1775
40	Leu Leu Gly Thr Ser Asn Ser Ala Ala Met Ser Met Glu Arg Asn Ile	1780	1785	1790
	Gly Thr Ile Asn Phe Lys Tyr Gly Gln Asp Gln Asn Thr Pro Arg Arg	1795	1800	1805
45	Phe Thr Leu Glu Gly Gly Ile Ala Gln Ala Asn Pro Gln Val Ala Ser	1810	1815	1820
	Ala Leu Thr Asp Leu Lys Lys Glu Gly Leu Glu Met Lys Ser	1825	1830	1835

This protein or polypeptide is about 198 kDa and has a pI of 8.98.

60 The present invention relates to an isolated DNA molecule having a nucleotide sequence of SEQ. ID. No. 9 as follows:

- 21 -

5 ATGACATCGT CACAGCAGCG GGTGAAAGG TTTTACAGT ATTTCTCCGC CGGGTGATAA 60  
 ACGCCCATAC ATCTGAAAGA CGGGGTGTGC GCCCTGTATA ACGAACAAGA TGAGGAGGCG 120  
 GCGGTGCTGG AAGTACCGCA ACACAGCGAC AGCCTGTTAC TACACTGCCG AATCATTGAG 180  
 GCTGACCCAC AAACCTCAAT AACCTGTAT TCGATGCTAT TACAGCTGAA TTTTGAAATG 240  
 10 GCGGCCATGC GCGGCTGTTG GCTGGCGCTG GATGAACTGC ACAACGTGCG TTTATGTTTT 300  
 CAGCAGTCGC TGGAGCATCT GGATGAAGCA AGTTTTAGCG ATATCGTTAG CGGCTTCATC 360  
 GAACATGCGG CAGAAGTGCG TGAGTATATA GCGCAATTAG ACGAGAGTAG CGCGGCATAA 420  
 15

This is known as the dspF gene. This isolated DNA molecule of the present invention encodes a hypersensitive response elicitor protein or polypeptide having an amino acid sequence of SEQ. ID. No. 10 as follows:

20 Met Thr Ser Ser Gln Gln Arg Val Glu Arg Phe Leu Gln Tyr Phe Ser  
 1 5 10 15  
 25 Ala Gly Cys Lys Thr Pro Ile His Leu Lys Asp Gly Val Cys Ala Leu  
 20 25 30  
 Tyr Asn Glu Gln Asp Glu Glu Ala Ala Val Leu Glu Val Pro Gln His  
 35 40 45  
 30 Ser Asp Ser Leu Leu Leu His Cys Arg Ile Ile Glu Ala Asp Pro Gln  
 50 55 60  
 Thr Ser Ile Thr Leu Tyr Ser Met Leu Leu Gln Leu Asn Phe Glu Met  
 65 70 75 80  
 35 Ala Ala Met Arg Gly Cys Trp Leu Ala Leu Asp Glu Leu His Asn Val  
 85 90 95  
 40 Arg Leu Cys Phe Gln Gln Ser Leu Glu His Leu Asp Glu Ala Ser Phe  
 100 105 110  
 Ser Asp Ile Val Ser Gly Phe Ile Glu His Ala Ala Glu Val Arg Glu  
 115 120 125  
 45 Tyr Ile Ala Gln Leu Asp Glu Ser Ser Ala Ala  
 130 135

This protein or polypeptide is about 16 kDa and has a pI of 4.45.

50 The hypersensitive response elicitor polypeptide or protein derived  
 from *Pseudomonas syringae* has an amino acid sequence corresponding to SEQ. ID.  
 No. 11 as follows:

55 Met Gln Ser Leu Ser Leu Asn Ser Ser Ser Leu Gln Thr Pro Ala Met  
 1 5 10 15

- 22 -

	Ala	Leu	Val	Leu	Val	Arg	Pro	Glu	Ala	Glu	Thr	Thr	Gly	Ser	Thr	Ser	
				20					25					30			
5	Ser	Lys	Ala	Leu	Gln	Glu	Val	Val	Val	Lys	Leu	Ala	Glu	Glu	Leu	Met	
			35					40					45				
	Arg	Asn	Gly	Gln	Leu	Asp	Asp	Ser	Ser	Pro	Leu	Gly	Lys	Leu	Leu	Ala	
		50					55					60					
	Lys	Ser	Met	Ala	Ala	Asp	Gly	Lys	Ala	Gly	Gly	Gly	Ile	Glu	Asp	Val	
	65					70					75					80	
10	Ile	Ala	Ala	Leu	Asp	Lys	Leu	Ile	His	Glu	Lys	Leu	Gly	Asp	Asn	Phe	
					85					90					95		
	Gly	Ala	Ser	Ala	Asp	Ser	Ala	Ser	Gly	Thr	Gly	Gln	Gln	Asp	Leu	Met	
				100					105					110			
15	Thr	Gln	Val	Leu	Asn	Gly	Leu	Ala	Lys	Ser	Met	Leu	Asp	Asp	Leu	Leu	
			115					120					125				
	Thr	Lys	Gln	Asp	Gly	Gly	Thr	Ser	Phe	Ser	Glu	Asp	Asp	Met	Pro	Met	
		130					135					140					
	Leu	Asn	Lys	Ile	Ala	Gln	Phe	Met	Asp	Asp	Asn	Pro	Ala	Gln	Phe	Pro	
	145					150					155					160	
20	Lys	Pro	Asp	Ser	Gly	Ser	Trp	Val	Asn	Glu	Leu	Lys	Glu	Asp	Asn	Phe	
					165					170					175		
	Leu	Asp	Gly	Asp	Glu	Thr	Ala	Ala	Phe	Arg	Ser	Ala	Leu	Asp	Ile	Ile	
				180					185					190			
25	Gly	Gln	Gln	Leu	Gly	Asn	Gln	Gln	Ser	Asp	Ala	Gly	Ser	Leu	Ala	Gly	
			195					200					205				
	Thr	Gly	Gly	Gly	Leu	Gly	Thr	Pro	Ser	Ser	Phe	Ser	Asn	Asn	Ser	Ser	
		210					215					220					
	Val	Met	Gly	Asp	Pro	Leu	Ile	Asp	Ala	Asn	Thr	Gly	Pro	Gly	Asp	Ser	
	225					230					235					240	
30	Gly	Asn	Thr	Arg	Gly	Glu	Ala	Gly	Gln	Leu	Ile	Gly	Glu	Leu	Ile	Asp	
					245					250					255		
	Arg	Gly	Leu	Gln	Ser	Val	Leu	Ala	Gly	Gly	Gly	Leu	Gly	Thr	Pro	Val	
				260					265					270			
35	Asn	Thr	Pro	Gln	Thr	Gly	Thr	Ser	Ala	Asn	Gly	Gly	Gln	Ser	Ala	Gln	
			275					280					285				
	Asp	Leu	Asp	Gln	Leu	Leu	Gly	Gly	Leu	Leu	Leu	Lys	Gly	Leu	Glu	Ala	
		290					295					300					
	Thr	Leu	Lys	Asp	Ala	Gly	Gln	Thr	Gly	Thr	Asp	Val	Gln	Ser	Ser	Ala	
	305					310					315					320	



- 23 -

Ala Gln Ile Ala Thr Leu Leu Val Ser Thr Leu Leu Gln Gly Thr Arg  
 325 330 335

Asn Gln Ala Ala Ala  
 340

5

This hypersensitive response elicitor polypeptide or protein has a molecular weight of 34-35 kDa. It is rich in glycine (about 13.5%) and lacks cysteine and tyrosine.

Further information about the hypersensitive response elicitor derived from

*Pseudomonas syringae* is found in He, S. Y., H. C. Huang, and A. Collmer,

10 "*Pseudomonas syringae* pv. *syringae* Harpin<sub>PSS</sub>: a Protein that is Secreted via the Hrp

Pathway and Elicits the Hypersensitive Response in Plants," Cell 73:1255-1266

(1993), which is hereby incorporated by reference. The DNA molecule encoding the hypersensitive response elicitor from *Pseudomonas syringae* has a nucleotide

sequence corresponding to SEQ. ID. No. 12 as follows:

15

ATGCAGAGTC TCAGTCTTAA CAGCAGCTCG CTGCAAACCC CGGCAATGGC CCTTGTCTCTG 60

GTACGTCTCTG AAGCCGAGAC GACTGGCAGT ACGTCGAGCA AGGCGCTTCA GGAAGTTGTCT 120

GTGAAGCTGG CCGAGGAACT GATGCGCAAT GGTCAACTCG ACGACAGCTC GCCATTGGGA 180

AAACTGTTGG CCAAGTCGAT GGCCGAGAT GGCAAGGCGG GCGGCGGTAT TGAGGATGTC 240

20

ATCGCTGCGC TGGACAAGCT GATCCATGAA AAGCTCGGTG ACAACTTCGG CGCGTCTGCG 300

GACAGCGCCT CGGGTACCGG ACAGCAGGAC CTGATGACTC AGGTGCTCAA TGGCCTGGCC 360

AAGTCGATGC TCGATGATCT TCTGACCAAG CAGGATGGCG GGACAAGCTT CTCCGAAGAC 420

GATATGCCGA TGCTGAACAA GATCGCGCAG TTCATGGATG ACAATCCCGC ACAGTTTCCC 480

AAGCCGGACT CGGGCTCCTG GGTGAACGAA CTCAAGGAAG ACAACTTCCT TGATGGCGAC 540

25

GAAACGGCTG CGTTCGGTTC GGCACCTCGAC ATCATTGGCC AGCAACTGGG TAATCAGCAG 600

AGTGACGCTG GCAGTCTGGC AGGGACGGGT GGAGGTCTGG GCACTCCGAG CAGTTTTTCC 660

AACAACTCGT CCGTGATGGG TGATCCGCTG ATCGACGCCA ATACCGGTCC CCGTGACAGC 720

GGCAATACCC GTGGTGAAGC GGGGCAACTG ATCGGCGAGC TTATCGACCG TGGCCTGCAA 780

TCGGTATTGG CCGGTGGTGG ACTGGGCACA CCCGTAAACA CCCCAGCAGC CCGTACGTCG 840

30

GCGAATGGCG GACAGTCCGC TCAGGATCTT GATCAGTTGC TGGGCGGCTT GCTGCTCAAG 900

GGCCTGGAGG CAACGCTCAA GGATGCCGGG CAAACAGGCA CCGACGTGCA GTCGAGCGCT 960

GCGCAAATCG CCACCTTGCT GGTCAGTACG CTGCTGCAAG GCACCCGCAA TCAGGCTGCA 1020

GCCTGA 1026

Another potentially suitable hypersensitive response elicitor from *Pseudomonas syringae* is disclosed in U.S. Patent Application Serial No. 09/120,817, which is hereby incorporated by reference. The protein has a nucleotide sequence of SEQ. ID. No. 13 as follows:

	TCCACTTCGC	TGATTTTGAA	ATTGGCAGAT	TCATAGAAAC	GTTCAGGTGT	GGAAATCAGG	60
	CTGAGTGC	AGATTTCGTT	GATAAGGGTG	TGGTACTGGT	CATTGTTGGT	CATTTCAAGG	120
10	CCTCTGAGTG	CGGTGCGGAG	CAATACCACT	CTTCCTGCTG	GCGTGTGCAC	ACTGAGTCGC	180
	AGGCATAGGC	ATTTCACTTC	CTTGCGTTGG	TTGGGCATAT	AAAAAAGGA	ACTTTTAAAA	240
15	ACAGTGCAAT	GAGATGCCGG	CAAAACGGGA	ACCGGTCGCT	GCGCTTTGCC	ACTCACTTCG	300
	AGCAAGCTCA	ACCCCAAACA	TCCACATCCC	TATCGAACGG	ACAGCGATAC	GGCCACTTGC	360
	TCTGGTAAAC	CCTGGAGCTG	GCGTCGGTCC	AATTGCCCAC	TTAGCGAGGT	AACGCAGCAT	420
20	GAGCATCGGC	ATCACACCCC	GGCCGCAACA	GACCACCACG	CCACTCGATT	TTTCGGCGCT	480
	AAGCGGCAAG	AGTCCTCAAC	CAAACACGTT	CGGCGAGCAG	AACACTCAGC	AAGCGATCGA	540
25	CCCGAGTGCA	CTGTTGTTCG	GCAGCGACAC	ACAGAAAGAC	GTCAACTTCG	GCACGCCCCGA	600
	CAGCACCGTC	CAGAATCCGC	AGGACGCCAG	CAAGCCCAAC	GACAGCCAGT	CCAACATCGC	660
	TAAATTGATC	AGTGCATTGA	TCATGTCGTT	GCTGCAGATG	CTCACCAACT	CCAATAAAAA	720
30	GCAGGACACC	AATCAGGAAC	AGCCTGATAG	CCAGGCTCCT	TTCCAGAACA	ACGGCGGGCT	780
	CGGTACACCG	TCGGCCGATA	GCGGGGGCGG	CGGTACACCG	GATGCGACAG	GTGGCGGGCGG	840
35	CGGTGATACG	CCAAGCGCAA	CAGGCGGTGG	CGGCGGTGAT	ACTCCGACCG	CAACAGGCGG	900
	TGGCGGCAGC	GGTGGCGGCG	GCACACCCAC	TGCAACAGGT	GGCGGCAGCG	GTGGCACACC	960
	CACTGCAACA	GGCGGTGGCG	AGGGTGGCGT	AACACCGCAA	ATCACTCCGC	AGTTGGCCAA	1020
40	CCCTAACCGT	ACCTCAGGTA	CTGGCTCGGT	GTCGGACACC	GCAGGTTCTA	CCGAGCAAGC	1080
	CGGCAAGATC	AATGTGGTGA	AAGACACCAT	CAAGGTCGGC	GCTGGCGAAG	TCTTTGACGG	1140
45	CCACGGCGCA	ACCTTCACTG	CCGACAAATC	TATGGGTAAC	GGAGACCAGG	GCGAAAATCA	1200
	GAAGCCCATG	TTGAGCTGG	CTGAAGGCGC	TACGTTGAAG	AATGTGAACC	TGGGTGAGAA	1260
	CGAGGTCGAT	GGCATCCACG	TGAAAGCCAA	AAACGCTCAG	GAAGTCACCA	TTGACAACGT	1320
50	GCATGCCCAG	AACGTCGGTG	AAGACCTGAT	TACGGTCAAA	GGCGAGGGAG	GCGCAGCGGT	1380
	CACTAATCTG	AACATCAAGA	ACAGCAGTGC	CAAAGGTGCA	GACGACAAGG	TTGTCCAGCT	1440
55	CAACGCCAAC	ACTCACTTGA	AAATCGACAA	CTTCAAGGCC	GACGATTTCT	GCACGATGGT	1500
	TCGCACCAAC	GGTGGCAAGC	AGTTTGATGA	CATGAGCATC	GAGCTGAACG	GCATCGAAGC	1560
60	TAACCACGGC	AAGTTGCCCC	TGGTGAAAAG	CGACAGTGAC	GATCTGAAGC	TGGCAACGGG	1620

- 25 -

CAACATCGCC ATGACCGACG TCAAACACGC CTACGATAAA ACCCAGGCAT CGACCCAACA 1680  
 CACCGAGCTT TGAATCCAGA CAAGTAGCTT GAAAAAAGGG GGTGGACTC 1729

5

This DNA molecule is known as the dspE gene for *Pseudomonas syringae*. This isolated DNA molecule of the present invention encodes a protein or polypeptide which elicits a plant pathogen's hypersensitive response having an amino acid sequence of SEQ. ID. No. 14 as follows:

10

Met Ser Ile Gly Ile Thr Pro Arg Pro Gln Gln Thr Thr Thr Pro Leu  
 1 5 10 15

15

Asp Phe Ser Ala Leu Ser Gly Lys Ser Pro Gln Pro Asn Thr Phe Gly  
 20 25 30

Glu Gln Asn Thr Gln Gln Ala Ile Asp Pro Ser Ala Leu Leu Phe Gly  
 35 40 45

20

Ser Asp Thr Gln Lys Asp Val Asn Phe Gly Thr Pro Asp Ser Thr Val  
 50 55 60

Gln Asn Pro Gln Asp Ala Ser Lys Pro Asn Asp Ser Gln Ser Asn Ile  
 65 70 75 80

25

Ala Lys Leu Ile Ser Ala Leu Ile Met Ser Leu Leu Gln Met Leu Thr  
 85 90 95

Asn Ser Asn Lys Lys Gln Asp Thr Asn Gln Glu Gln Pro Asp Ser Gln  
 100 105 110

30

Ala Pro Phe Gln Asn Asn Gly Gly Leu Gly Thr Pro Ser Ala Asp Ser  
 115 120 125

Gly Gly Gly Gly Thr Pro Asp Ala Thr Gly Gly Gly Gly Gly Asp Thr  
 130 135 140

35

Pro Ser Ala Thr Gly Gly Gly Gly Gly Asp Thr Pro Thr Ala Thr Gly  
 145 150 155 160

40

Gly Gly Gly Ser Gly Gly Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly  
 165 170 175

Ser Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly Glu Gly Gly Val Thr  
 180 185 190

45

Pro Gln Ile Thr Pro Gln Leu Ala Asn Pro Asn Arg Thr Ser Gly Thr  
 195 200 205

Gly Ser Val Ser Asp Thr Ala Gly Ser Thr Glu Gln Ala Gly Lys Ile  
 210 215 220

50

Asn Val Val Lys Asp Thr Ile Lys Val Gly Ala Gly Glu Val Phe Asp  
 225 230 235 240

55

- 26 -

	Gly	His	Gly	Ala	Thr	Phe	Thr	Ala	Asp	Lys	Ser	Met	Gly	Asn	Gly	Asp	
					245					250					255		
5	Gln	Gly	Glu	Asn	Gln	Lys	Pro	Met	Phe	Glu	Leu	Ala	Glu	Gly	Ala	Thr	
				260					265						270		
	Leu	Lys	Asn	Val	Asn	Leu	Gly	Glu	Asn	Glu	Val	Asp	Gly	Ile	His	Val	
			275					280					285				
10	Lys	Ala	Lys	Asn	Ala	Gln	Glu	Val	Thr	Ile	Asp	Asn	Val	His	Ala	Gln	
		290					295					300					
	Asn	Val	Gly	Glu	Asp	Leu	Ile	Thr	Val	Lys	Gly	Glu	Gly	Gly	Ala	Ala	
15		305				310					315					320	
	Val	Thr	Asn	Leu	Asn	Ile	Lys	Asn	Ser	Ser	Ala	Lys	Gly	Ala	Asp	Asp	
				325					330						335		
	Lys	Val	Val	Gln	Leu	Asn	Ala	Asn	Thr	His	Leu	Lys	Ile	Asp	Asn	Phe	
20				340					345					350			
	Lys	Ala	Asp	Asp	Phe	Gly	Thr	Met	Val	Arg	Thr	Asn	Gly	Gly	Lys	Gln	
			355					360					365				
25	Phe	Asp	Asp	Met	Ser	Ile	Glu	Leu	Asn	Gly	Ile	Glu	Ala	Asn	His	Gly	
		370					375					380					
	Lys	Phe	Ala	Leu	Val	Lys	Ser	Asp	Ser	Asp	Asp	Leu	Lys	Leu	Ala	Thr	
30		385				390				395						400	
	Gly	Asn	Ile	Ala	Met	Thr	Asp	Val	Lys	His	Ala	Tyr	Asp	Lys	Thr	Gln	
					405				410						415		
35	Ala	Ser	Thr	Gln	His	Thr	Glu	Leu									
				420													

This protein or polypeptide is about 42.9 kDa.

40                   The hypersensitive response elicitor polypeptide or protein derived from *Pseudomonas solanacearum* has an amino acid sequence corresponding to SEQ. ID. No. 15 as follows:

45	Met	Ser	Val	Gly	Asn	Ile	Gln	Ser	Pro	Ser	Asn	Leu	Pro	Gly	Leu	Gln	
	1				5					10					15		
	Asn	Leu	Asn	Leu	Asn	Thr	Asn	Thr	Asn	Ser	Gln	Gln	Ser	Gly	Gln	Ser	
				20					25					30			
	Val	Gln	Asp	Leu	Ile	Lys	Gln	Val	Glu	Lys	Asp	Ile	Leu	Asn	Ile	Ile	
			35				40					45					
50	Ala	Ala	Leu	Val	Gln	Lys	Ala	Ala	Gln	Ser	Ala	Gly	Gly	Asn	Thr	Gly	
		50				55						60					

- 27 -

	Asn Thr Gly Asn Ala Pro Ala Lys Asp Gly Asn Ala Asn Ala Gly Ala	65	70	75	80
	Asn Asp Pro Ser Lys Asn Asp Pro Ser Lys Ser Gln Ala Pro Gln Ser	85	90	95	
5	Ala Asn Lys Thr Gly Asn Val Asp Asp Ala Asn Asn Gln Asp Pro Met	100	105	110	
	Gln Ala Leu Met Gln Leu Leu Glu Asp Leu Val Lys Leu Leu Lys Ala	115	120	125	
10	Ala Leu His Met Gln Gln Pro Gly Gly Asn Asp Lys Gly Asn Gly Val	130	135	140	
	Gly Gly Ala Asn Gly Ala Lys Gly Ala Gly Gly Gln Gly Gly Leu Ala	145	150	155	160
	Glu Ala Leu Gln Glu Ile Glu Gln Ile Leu Ala Gln Leu Gly Gly Gly	165	170	175	
15	Gly Ala Gly Ala Gly Gly Ala Gly Gly Gly Val Gly Gly Ala Gly Gly	180	185	190	
	Ala Asp Gly Gly Ser Gly Ala Gly Gly Ala Gly Gly Ala Asn Gly Ala	195	200	205	
20	Asp Gly Gly Asn Gly Val Asn Gly Asn Gln Ala Asn Gly Pro Gln Asn	210	215	220	
	Ala Gly Asp Val Asn Gly Ala Asn Gly Ala Asp Asp Gly Ser Glu Asp	225	230	235	240
	Gln Gly Gly Leu Thr Gly Val Leu Gln Lys Leu Met Lys Ile Leu Asn	245	250	255	
25	Ala Leu Val Gln Met Met Gln Gln Gly Gly Leu Gly Gly Gly Asn Gln	260	265	270	
	Ala Gln Gly Gly Ser Lys Gly Ala Gly Asn Ala Ser Pro Ala Ser Gly	275	280	285	
30	Ala Asn Pro Gly Ala Asn Gln Pro Gly Ser Ala Asp Asp Gln Ser Ser	290	295	300	
	Gly Gln Asn Asn Leu Gln Ser Gln Ile Met Asp Val Val Lys Glu Val	305	310	315	320
	Val Gln Ile Leu Gln Gln Met Leu Ala Ala Gln Asn Gly Gly Ser Gln	325	330	335	
35	Gln Ser Thr Ser Thr Gln Pro Met	340			

It is encoded by a DNA molecule having a nucleotide sequence corresponding SEQ.  
ID. No. 16 as follows:

- 28 -

ATGTCAGTCG GAAACATCCA GAGCCCGTCG AACCTCCCGG GTCTGCAGAA CCTGAACCTC 60  
 AACACCAACA CCAACAGCCA GCAATCGGGC CAGTCCGTGC AAGACCTGAT CAAGCAGGTC 120  
 GAGAAGGACA TCCTCAACAT CATCGCAGCC CTCGTGCAGA AGGCCGCACA GTCGGCGGGC 180  
 GGCAACACCG GTAACACCGG CAACGCGCCG GCGAAGGACG GCAATGCCAA CGCGGGCGCC 240  
 5 AACGACCCGA GCAAGAACGA CCCGAGCAAG AGCCAGGCTC CGCAGTCGGC CAACAAGACC 300  
 GGCAACGTCG ACGACGCCAA CAACCAGGAT CCGATGCAAG CGCTGATGCA GCTGCTGGAA 360  
 GACCTGGTGA AGCTGCTGAA GGCGGCCCTG CACATGCAGC AGCCCGGCGG CAATGACAAG 420  
 GGCAACGGCG TGGGCGGTGC CAACGGCGCC AAGGGTGCCG GCGGCCAGGG CGGCCTGGCC 480  
 GAAGCGCTGC AGGAGATCGA GCAGATCCTC GCCCAGCTCG GCGGCGGCGG TGCTGGCGCC 540  
 10 GGCGGCGCGG GTGGCGGTGT CGGCGGTGCT GGTGGCGCGG ATGGCGGCTC CGGTGCGGGT 600  
 GGCGCAGGCG GTGCGAACGG CGCCGACGGC GGCAATGGCG TGAACGGCAA CCAGGCGAAC 660  
 GGCCCGCAGA ACGCAGGCGA TGTCAACGGT GCCAACGGCG CGGATGACGG CAGCGAAGAC 720  
 CAGGCGCGCC TCACCGGCGT GCTGCAAAAG CTGATGAAGA TCCTGAACGC GCTGGTGCAG 780  
 ATGATGCAGC AAGGCGGCCT CGGCGGCGGC AACCAGGCGC AGGGCGGCTC GAAGGGTGCC 840  
 15 GGCAACGCCT CGCCGGCTTC CGGCGCGAAC CCGGGCGCGA ACCAGCCCGG TTCGGCGGAT 900  
 GATCAATCGT CCGGCCAGAA CAATCTGCAA TCCCAGATCA TGGATGTGGT GAAGGAGGTC 960  
 GTCCAGATCC TGCAGCAGAT GCTGGCGGCG CAGAACGGCG GCAGCCAGCA GTCCACCTCG 1020  
 ACGCAGCCGA TGTA 1035

20

Further information regarding the hypersensitive response elicitor polypeptide or protein derived from *Pseudomonas solanacearum* is set forth in Arlat, M., F. Van Gijsegem, J. C. Huet, J. C. Pemollet, and C. A. Boucher, "PopA1, a Protein which Induces a Hypersensitive-like Response in Specific Petunia Genotypes, is Secreted via the Hrp Pathway of *Pseudomonas solanacearum*," EMBO J. 13:543-533 (1994), which is hereby incorporated by reference.

25

The hypersensitive response elicitor polypeptide or protein from *Xanthomonas campestris* pv. *glycines* has an amino acid sequence corresponding to SEQ. ID. No. 17 as follows:

30

Thr Leu Ile Glu Leu Met Ile Val Val Ala Ile Ile Ala Ile Leu Ala  
 1 5 10 15

- 29 -

Ala Ile Ala Leu Pro Ala Tyr Gln Asp Tyr  
20 25

5 This sequence is an amino terminal sequence having only 26 residues from the hypersensitive response elicitor polypeptide or protein of *Xanthomonas campestris* pv. glycines. It matches with fimbrial subunit proteins determined in other *Xanthomonas campestris* pathovars.

The hypersensitive response elicitor polypeptide or protein from  
10 *Xanthomonas campestris* pv. *pelargonii* is heat stable, protease sensitive, and has a molecular weight of 20 kDa. It includes an amino acid sequence corresponding to SEQ. ID. No. 18 as follows:

15 Ser Ser Gln Gln Ser Pro Ser Ala Gly Ser Glu Gln Gln Leu Asp Gln  
1 5 10 15  
Leu Leu Ala Met  
20

20 Isolation of *Erwinia carotovora* hypersensitive response elicitor protein or polypeptide is described in Cui et al., "The RsmA Mutants of *Erwinia carotovora* subsp. *carotovora* Strain Ecc71 Overexpress *hrp N<sub>Ecc</sub>* and Elicit a Hypersensitive Reaction-like Response in Tobacco Leaves," MPMI, 9(7):565-73 (1996), which is hereby incorporated by reference. The hypersensitive response elicitor protein or  
25 polypeptide of *Erwinia stewartii* is set forth in Ahmad et al., "Harpin is Not Necessary for the Pathogenicity of *Erwinia stewartii* on Maize," 8th Int'l. Cong. Molec. Plant-Microbe Interact., July 14-19, 1996 and Ahmad, et al., "Harpin is Not Necessary for the Pathogenicity of *Erwinia stewartii* on Maize," Ann. Mtg. Am. Phytopath. Soc., July 27-31, 1996, which are hereby incorporated by reference.

30 Hypersensitive response elicitor proteins or polypeptides from *Phytophthora parasitica*, *Phytophthora cryptogea*, *Phytophthora cinnamoni*, *Phytophthora capsici*, *Phytophthora megasperma*, and *Phytophthora citrophthora* are described in Kaman, et al., "Extracellular Protein Elicitors from *Phytophthora*: Most Specificity and Induction of Resistance to Bacterial and Fungal Phytopathogens,"  
35 Molec. Plant-Microbe Interact., 6(1):15-25 (1993), Ricci et al., "Structure and Activity of Proteins from Pathogenic Fungi *Phytophthora* Eliciting Necrosis and

Acquired Resistance in Tobacco," Eur. J. Biochem., 183:555-63 (1989), Ricci et al., "Differential Production of Parasiticein, and Elicitor of Necrosis and Resistance in Tobacco, by Isolates of *Phytophthora parasitica*," Plant Path. 41:298-307 (1992), Baillreul et al, "A New Elicitor of the Hypersensitive Response in Tobacco: A Fungal Glycoprotein Elicits Cell Death, Expression of Defence Genes, Production of Salicylic Acid, and Induction of Systemic Acquired Resistance," Plant J., 8(4):551-60 (1995), and Bonnet et al., "Acquired Resistance Triggered by Elicitors in Tobacco and Other Plants," Eur. J. Plant Path., 102:181-92 (1996), which are hereby incorporated by reference.

Another hypersensitive response elicitor in accordance with the present invention is from *Clavibacter michiganensis* subsp. *sepedonicus* which is fully described in U.S. Patent Application Serial No. 09/136,625, which is hereby incorporated by reference.

The above elicitors are exemplary. Other elicitors can be identified by growing fungi or bacteria that elicit a hypersensitive response under conditions which genes encoding an elicitor are expressed. Cell-free preparations from culture supernatants can be tested for elicitor activity (i.e. local necrosis) by using them to infiltrate appropriate plant tissues.

Fragments of the above hypersensitive response elicitor polypeptides or proteins as well as fragments of full length elicitors from other pathogens are encompassed by the method of the present invention.

Suitable fragments can be produced by several means. In the first, subclones of the gene encoding a known elicitor protein are produced by conventional molecular genetic manipulation by subcloning gene fragments. The subclones then are expressed *in vitro* or *in vivo* in bacterial cells to yield a smaller protein or peptide that can be tested for elicitor activity according to the procedure described below.

As an alternative, fragments of an elicitor protein can be produced by digestion of a full-length elicitor protein with proteolytic enzymes like chymotrypsin or *Staphylococcus* proteinase A, or trypsin. Different proteolytic enzymes are likely to cleave elicitor proteins at different sites based on the amino acid sequence of the elicitor protein. Some of the fragments that result from proteolysis may be active elicitors of resistance.



In another approach, based on knowledge of the primary structure of the protein, fragments of the elicitor protein gene may be synthesized by using the PCR technique together with specific sets of primers chosen to represent particular portions of the protein. These then would be cloned into an appropriate vector for  
5 expression of a truncated peptide or protein.

Chemical synthesis can also be used to make suitable fragments. Such a synthesis is carried out using known amino acid sequences for the elicitor being produced. Alternatively, subjecting a full length elicitor to high temperatures and pressures will produce fragments. These fragments can then be separated by  
10 conventional procedures (e.g., chromatography, SDS-PAGE).

An example of suitable fragments of a hypersensitive-response elicitor which do not elicit a hypersensitive response include fragments of the *Erwinia*. Suitable fragments include a C-terminal fragment of the amino acid sequence of SEQ. ID. No. 3, an N-terminal fragment of the amino acid sequence of SEQ. ID. No. 3, or  
15 an internal fragment of the amino acid sequence of SEQ. ID. No. 3. The C-terminal fragment of the amino acid sequence of SEQ. ID. No. 3 can span the following amino acids of SEQ. ID. No. 3: 169 and 403, 210 and 403, 267 and 403, or 343 and 403. The internal fragment of the amino acid sequence of SEQ. ID. No. 3 can span the following amino acids of SEQ. ID. No. 3: 105 and 179, 137 and 166, 121 and 150, or  
20 137 and 156. Other suitable fragments can be identified in accordance with the present invention.

Another example of suitable fragments of a hypersensitive response elicitor which do elicit a hypersensitive response are *Erwinia amylovora* fragments including a C-terminal fragment of the amino acid sequence of SEQ. ID. No. 3, an N-  
25 terminal fragment of the amino acid sequence of SEQ. ID. No. 3, or an internal fragment of the amino acid sequence of SEQ. ID. No. 3. The C-terminal fragment of the amino acid sequence of SEQ. ID. No. 3 can span amino acids 105 and 403 of SEQ. ID. No. 3. The N-terminal fragment of the amino acid sequence of SEQ. ID. No. 3 can span the following amino acids of SEQ. ID. No. 3: 1 and 98, 1 and 104, 1  
30 and 122, 1 and 168, 1 and 218, 1 and 266, 1 and 342, 1 and 321, and 1 and 372. The internal fragment of the amino acid sequence of SEQ. ID. No. 3 can span the

- 32 -

following amino acids of SEQ. ID. No. 3: 76 and 209, 105 and 209, 99 and 209, 137 and 204, 137 and 200, 109 and 204, 109 and 200, 137 and 180, and 105 and 180.

Suitable DNA molecules are those that hybridize to the DNA molecule comprising a nucleotide sequence of SEQ. ID. Nos. 2, 4, 5, 7, 9, 12, 13, and 16 under  
5 stringent conditions. An example of suitable high stringency conditions is when hybridization is carried out at 65°C for 20 hours in a medium containing 1M NaCl, 50 mM Tris-HCl, pH 7.4, 10 mM EDTA, 0.1% sodium dodecyl sulfate, 0.2% ficoll, 0.2% polyvinylpyrrolidone, 0.2% bovine serum albumin, 50 µm g/ml *E. coli* DNA.

Variants may be made by, for example, the deletion or addition of  
10 amino acids that have minimal influence on the properties, secondary structure and hydropathic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis,  
15 purification, or identification of the polypeptide.

The hypersensitive response elicitor of the present invention is preferably in isolated form (i.e. separated from its host organism) and more preferably produced in purified form (preferably at least about 60%, more preferably 80%, pure) by conventional techniques. Typically, the hypersensitive response elicitor of the  
20 present invention is produced but not secreted into the growth medium of recombinant host cells. Alternatively, the protein or polypeptide of the present invention is secreted into growth medium. In the case of unsecreted protein, to isolate the protein, the host cell (e.g., *E. coli*) carrying a recombinant plasmid is propagated, lysed by sonication, heat, or chemical treatment, and the homogenate is centrifuged to remove  
25 bacterial debris. The supernatant is then subjected to heat treatment and the hypersensitive response elicitor is separated by centrifugation. The supernatant fraction containing the hypersensitive response elicitor is subjected to gel filtration in an appropriately sized dextran or polyacrylamide column to separate the fragment. If necessary, the protein fraction may be further purified by ion exchange or HPLC.

30 The DNA molecule encoding the hypersensitive response elicitor polypeptide or protein can be incorporated in cells using conventional recombinant DNA technology. Generally, this involves inserting the DNA molecule into an

- 33 -

expression system to which the DNA molecule is heterologous (i.e. not normally present). The heterologous DNA molecule is inserted into the expression system or vector in sense orientation and correct reading frame. The vector contains the necessary elements for the transcription and translation of the inserted protein-coding sequences.

U.S. Patent No. 4,237,224 to Cohen and Boyer, which is hereby incorporated by reference, describes the production of expression systems in the form of recombinant plasmids using restriction enzyme cleavage and ligation with DNA ligase. These recombinant plasmids are then introduced by means of transformation and replicated in unicellular cultures including procaryotic organisms and eucaryotic cells grown in tissue culture.

Recombinant genes may also be introduced into viruses, such as vaccina virus. Recombinant viruses can be generated by transfection of plasmids into cells infected with virus.

Suitable vectors include, but are not limited to, the following viral vectors such as lambda vector system gt11, gt WES.tB, Charon 4, and plasmid vectors such as pBR322, pBR325, pACYC177, pACYC1084, pUC8, pUC9, pUC18, pUC19, pLG339, pR290, pKC37, pKC101, SV 40, pBluescript II SK +/- or KS +/- (see "Stratagene Cloning Systems" Catalog (1993) from Stratagene, La Jolla, Calif, which is hereby incorporated by reference), pQE, pIH821, pGEX, pET series (see F.W. Studier et. al., "Use of T7 RNA Polymerase to Direct Expression of Cloned Genes," Gene Expression Technology vol. 185 (1990), which is hereby incorporated by reference), and any derivatives thereof. Recombinant molecules can be introduced into cells via transformation, particularly transduction, conjugation, mobilization, or electroporation. The DNA sequences are cloned into the vector using standard cloning procedures in the art, as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Springs Laboratory, Cold Springs Harbor, New York (1989), which is hereby incorporated by reference.

A variety of host-vector systems may be utilized to express the protein-encoding sequence(s). Primarily, the vector system must be compatible with the host cell used. Host-vector systems include but are not limited to the following: bacteria transformed with bacteriophage DNA, plasmid DNA, or cosmid DNA;

- 34 -

microorganisms such as yeast containing yeast vectors; mammalian cell systems infected with virus (e.g., vaccinia virus, adenovirus, etc.); insect cell systems infected with virus (e.g., baculovirus); and plant cells infected by bacteria. The expression elements of these vectors vary in their strength and specificities. Depending upon the host-vector system utilized, any one of a number of suitable transcription and translation elements can be used.

Different genetic signals and processing events control many levels of gene expression (e.g., DNA transcription and messenger RNA (mRNA) translation).

Transcription of DNA is dependent upon the presence of a promotor which is a DNA sequence that directs the binding of RNA polymerase and thereby promotes mRNA synthesis. The DNA sequences of eucaryotic promotors differ from those of procaryotic promotors. Furthermore, eucaryotic promotors and accompanying genetic signals may not be recognized in or may not function in a procaryotic system, and, further, procaryotic promotors are not recognized and do not function in eucaryotic cells.

Similarly, translation of mRNA in procaryotes depends upon the presence of the proper procaryotic signals which differ from those of eucaryotes. Efficient translation of mRNA in procaryotes requires a ribosome binding site called the Shine-Dalgarno ("SD") sequence on the mRNA. This sequence is a short nucleotide sequence of mRNA that is located before the start codon, usually AUG, which encodes the amino-terminal methionine of the protein. The SD sequences are complementary to the 3'-end of the 16S rRNA (ribosomal RNA) and probably promote binding of mRNA to ribosomes by duplexing with the rRNA to allow correct positioning of the ribosome. For a review on maximizing gene expression, see Roberts and Lauer, Methods in Enzymology, 68:473 (1979), which is hereby incorporated by reference.

Promotors vary in their "strength" (i.e. their ability to promote transcription). For the purposes of expressing a cloned gene, it is desirable to use strong promotors in order to obtain a high level of transcription and, hence, expression of the gene. Depending upon the host cell system utilized, any one of a number of suitable promotors may be used. For instance, when cloning in *E. coli*, its bacteriophages, or plasmids, promotors such as the T7 phage promotor, *lac* promotor,

*trp* promotor, *recA* promotor, ribosomal RNA promotor, the  $P_R$  and  $P_L$  promoters of coliphage lambda and others, including but not limited, to *lacUV5*, *ompF*, *bla*, *lpp*, and the like, may be used to direct high levels of transcription of adjacent DNA segments. Additionally, a hybrid *trp-lacUV5* (*tac*) promotor or other *E. coli* 5 promoters produced by recombinant DNA or other synthetic DNA techniques may be used to provide for transcription of the inserted gene.

Bacterial host cell strains and expression vectors may be chosen which inhibit the action of the promotor unless specifically induced. In certain operations, the addition of specific inducers is necessary for efficient transcription of the inserted 10 DNA. For example, the *lac* operon is induced by the addition of lactose or IPTG (isopropylthio-beta-D-galactoside). A variety of other operons, such as *trp*, *pro*, etc., are under different controls.

Specific initiation signals are also required for efficient gene transcription and translation in procaryotic cells. These transcription and translation 15 initiation signals may vary in "strength" as measured by the quantity of gene specific messenger RNA and protein synthesized, respectively. The DNA expression vector, which contains a promotor, may also contain any combination of various "strong" transcription and/or translation initiation signals. For instance, efficient translation in *E. coli* requires an SD sequence about 7-9 bases 5' to the initiation codon ("ATG") to 20 provide a ribosome binding site. Thus, any SD-ATG combination that can be utilized by host cell ribosomes may be employed. Such combinations include but are not limited to the SD-ATG combination from the *cro* gene or the *N* gene of coliphage lambda, or from the *E. coli* tryptophan E, D, C, B or A genes. Additionally, any SD-ATG combination produced by recombinant DNA or other techniques involving 25 incorporation of synthetic nucleotides may be used.

Once the isolated DNA molecule encoding the hypersensitive response elicitor polypeptide or protein has been cloned into an expression system, it is ready to be incorporated into a host cell. Such incorporation can be carried out by the various forms of transformation noted above, depending upon the vector/host cell 30 system. Suitable host cells include, but are not limited to, bacteria, virus, yeast, mammalian cells, insect, plant, and the like.

- 36 -

The present invention's method of imparting stress resistance to plants can involve applying the hypersensitive response elicitor polypeptide or protein in a non-infectious form to all or part of a plant or a plant seed under conditions effective for the elicitor to impart stress resistance. Alternatively, the hypersensitive response elicitor protein or polypeptide can be applied to plants such that seeds recovered from such plants themselves are able to impart stress resistance in plants.

As an alternative to applying a hypersensitive response elicitor polypeptide or protein to plants or plant seeds in order to impart stress resistance in plants or plants grown from the seeds, transgenic plants or plant seeds can be utilized. When utilizing transgenic plants, this involves providing a transgenic plant transformed with a DNA molecule encoding a hypersensitive response elicitor polypeptide or protein and growing the plant under conditions effective to permit that DNA molecule to impart stress resistance to plants. Alternatively, a transgenic plant seed transformed with a DNA molecule encoding a hypersensitive response elicitor polypeptide or protein can be provided and planted in soil. A plant is then propagated from the planted seed under conditions effective to permit that DNA molecule to impart stress resistance to plants.

The embodiment of the present invention where the hypersensitive response elicitor-polypeptide or protein is applied to the plant or plant seed can be carried out in a number of ways, including: 1) application of an isolated hypersensitive response elicitor or 2) application of bacteria which do not cause disease and are transformed with a genes encoding the elicitor. In the latter embodiment, the elicitor can be applied to plants or plant seeds by applying bacteria containing the DNA molecule encoding a hypersensitive response elicitor polypeptide or protein. Such bacteria must be capable of secreting or exporting the elicitor so that the elicitor can contact plant or plant seed cells. In these embodiments, the elicitor is produced by the bacteria *in planta* or on seeds or just prior to introduction of the bacteria to the plants or plant seeds.

The methods of the present invention can be utilized to treat a wide variety of plants or their seeds to impart stress resistance. Suitable plants include dicots and monocots. More particularly, useful crop plants can include: alfalfa, rice, wheat, barley, rye, cotton, sunflower, peanut, corn, potato, sweet potato, bean, pea,

chicory, lettuce, endive, cabbage, brussel sprout, beet, parsnip, cauliflower, broccoli, turnip, radish, spinach, onion, garlic, eggplant, pepper, celery, carrot, squash, pumpkin, zucchini, cucumber, apple, pear, melon, citrus, strawberry, grape, raspberry, pineapple, soybean, tobacco, tomato, sorghum, and sugarcane. Examples of suitable  
5 ornamental plants are: *Arabidopsis thaliana*, *Saintpaulia*, petunia, pelargonium, poinsettia, chrysanthemum, carnation, and zinnia.

In accordance with the present invention, the term "stress" refers to drought, salt, cold temperatures (e.g., frost), chemical treatment (e.g., insecticides, fungicides, herbicides, fertilizers), water, excessive light, and insufficient light.

10 The method of the present invention involving application of the hypersensitive response elicitor polypeptide or protein can be carried out through a variety of procedures when all or part of the plant is treated, including leaves, stems, roots, propagules (e.g., cuttings), etc. This may (but need not) involve infiltration of the hypersensitive response elicitor polypeptide or protein into the plant. Suitable  
15 application methods include high or low pressure spraying, injection, and leaf abrasion proximate to when elicitor application takes place. When treating plant seeds or propagules (e.g., cuttings), in accordance with the application embodiment of the present invention, the hypersensitive response elicitor protein or polypeptide, in accordance with present invention, can be applied by low or high pressure spraying,  
20 coating, immersion, or injection. Other suitable application procedures can be envisioned by those skilled in the art provided they are able to effect contact of the elicitor with cells of the plant or plant seed. Once treated with the hypersensitive response elicitor of the present invention, the seeds can be planted in natural or artificial soil and cultivated using conventional procedures to produce plants. After  
25 plants have been propagated from seeds treated in accordance with the present invention, the plants may be treated with one or more applications of the hypersensitive response elicitor protein or polypeptide to impart stress resistance to plants.

The hypersensitive response elicitor polypeptide or protein, in  
30 accordance with the present invention, can be applied to plants or plant seeds alone or in a mixture with other materials. Alternatively, the hypersensitive response elicitor

- 38 -

polypeptide or protein can be applied separately to plants with other materials being applied at different times.

A composition suitable for treating plants or plant seeds in accordance with the application embodiment of the present invention contains a hypersensitive response elicitor polypeptide or protein in a carrier. Suitable carriers include water, aqueous solutions, slurries, or dry powders. In this embodiment, the composition contains greater than 500 nM of the elicitor.

Although not required, this composition may contain additional additives including fertilizer, insecticide, fungicide, nematocide, and mixtures thereof. Suitable fertilizers include  $(\text{NH}_4)_2\text{NO}_3$ . An example of a suitable insecticide is Malathion. Useful fungicides include Captan.

Other suitable additives include buffering agents, wetting agents, coating agents, and abrading agents. These materials can be used to facilitate the process of the present invention. In addition, the hypersensitive response elicitor can be applied to plant seeds with other conventional seed formulation and treatment materials, including clays and polysaccharides.

In the alternative embodiment of the present invention involving the use of transgenic plants and transgenic seeds, a hypersensitive response elicitor need not be applied topically to the plants or seeds. Instead, transgenic plants transformed with a DNA molecule encoding such an elicitor are produced according to procedures well known in the art.

The vector described above can be microinjected directly into plant cells by use of micropipettes to transfer mechanically the recombinant DNA. Crossway, Mol. Gen. Genetics, 202:179-85 (1985), which is hereby incorporated by reference. The genetic material may also be transferred into the plant cell using polyethylene glycol. Krens, et al., Nature, 296:72-74 (1982), which is hereby incorporated by reference.

Another approach to transforming plant cells with a gene is particle bombardment (also known as biolistic transformation) of the host cell. This can be accomplished in one of several ways. The first involves propelling inert or biologically active particles at cells. This technique is disclosed in U.S. Patent Nos. 4,945,050, 5,036,006, and 5,100,792, all to Sanford et al., which are hereby



incorporated by reference. Generally, this procedure involves propelling inert or biologically active particles at the cells under conditions effective to penetrate the outer surface of the cell and to be incorporated within the interior thereof. When inert particles are utilized, the vector can be introduced into the cell by coating the particles with the vector containing the heterologous DNA. Alternatively, the target cell can be surrounded by the vector so that the vector is carried into the cell by the wake of the particle. Biologically active particles (e.g., dried bacterial cells containing the vector and heterologous DNA) can also be propelled into plant cells.

Yet another method of introduction is fusion of protoplasts with other entities, either minicells, cells, lysosomes, or other fusible lipid-surfaced bodies. Fraley, et al., Proc. Natl. Acad. Sci. USA, 79:1859-63 (1982), which is hereby incorporated by reference.

The DNA molecule may also be introduced into the plant cells by electroporation. Fromm et al., Proc. Natl. Acad. Sci. USA, 82:5824 (1985), which is hereby incorporated by reference. In this technique, plant protoplasts are electroporated in the presence of plasmids containing the expression cassette. Electrical impulses of high field strength reversibly permeabilize biomembranes allowing the introduction of the plasmids. Electroporated plant protoplasts reform the cell wall, divide, and regenerate.

Another method of introducing the DNA molecule into plant cells is to infect a plant cell with *Agrobacterium tumefaciens* or *A. rhizogenes* previously transformed with the gene. Under appropriate conditions known in the art, the transformed plant cells are grown to form shoots or roots, and develop further into plants. Generally, this procedure involves inoculating the plant tissue with a suspension of bacteria and incubating the tissue for 48 to 72 hours on regeneration medium without antibiotics at 25-28°C.

*Agrobacterium* is a representative genus of the Gram-negative family Rhizobiaceae. Its species are responsible for crown gall (*A. tumefaciens*) and hairy root disease (*A. rhizogenes*). The plant cells in crown gall tumors and hairy roots are induced to produce amino acid derivatives known as opines, which are catabolized only by the bacteria. The bacterial genes responsible for expression of opines are a

convenient source of control elements for chimeric expression cassettes. In addition, assaying for the presence of opines can be used to identify transformed tissue.

Heterologous genetic sequences can be introduced into appropriate plant cells, by means of the Ti plasmid of *A. tumefaciens* or the Ri plasmid of *A. rhizogenes*. The Ti or Ri plasmid is transmitted to plant cells on infection by Agrobacterium and is stably integrated into the plant genome. J. Schell, Science, 237:1176-83 (1987), which is hereby incorporated by reference.

After transformation, the transformed plant cells must be regenerated.

Plant regeneration from cultured protoplasts is described in Evans et al., Handbook of Plant Cell Cultures, Vol. 1: (MacMillan Publishing Co., New York, 1983); and Vasil I.R. (ed.), Cell Culture and Somatic Cell Genetics of Plants, Acad. Press, Orlando, Vol. I, 1984, and Vol. III (1986), which are hereby incorporated by reference.

It is known that practically all plants can be regenerated from cultured cells or tissues, including but not limited to, all major species of sugarcane, sugar beets, cotton, fruit trees, and legumes.

Means for regeneration vary from species to species of plants, but generally a suspension of transformed protoplasts or a petri plate containing transformed explants is first provided. Callus tissue is formed and shoots may be induced from callus and subsequently rooted. Alternatively, embryo formation can be induced in the callus tissue. These embryos germinate as natural embryos to form plants. The culture media will generally contain various amino acids and hormones, such as auxin and cytokinins. It is also advantageous to add glutamic acid and proline to the medium, especially for such species as corn and alfalfa. Efficient regeneration will depend on the medium, on the genotype, and on the history of the culture. If these three variables are controlled, then regeneration is usually reproducible and repeatable.

After the expression cassette is stably incorporated in transgenic plants, it can be transferred to other plants by sexual crossing. Any of a number of standard breeding techniques can be used, depending upon the species to be crossed.

Once transgenic plants of this type are produced, the plants themselves can be cultivated in accordance with conventional procedure with the presence of the

gene encoding the hypersensitive response elicitor resulting in stress resistance to the plant. Alternatively, transgenic seeds or propagules (e.g., cuttings) are recovered from the transgenic plants. The seeds can then be planted in the soil and cultivated using conventional procedures to produce transgenic plants. The transgenic plants are propagated from the planted transgenic seeds under conditions effective to impart stress resistance to plants. While not wishing to be bound by theory, such stress resistance may be RNA mediated or may result from expression of the elicitor polypeptide or protein.

When transgenic plants and plant seeds are used in accordance with the present invention, they additionally can be treated with the same materials as are used to treat the plants and seeds to which a hypersensitive response elicitor in accordance with the present invention is applied. These other materials, including a hypersensitive response elicitor in accordance with the present invention, can be applied to the transgenic plants and plant seeds by the above-noted procedures, including high or low pressure spraying, injection, coating, and immersion. Similarly, after plants have been propagated from the transgenic plant seeds, the plants may be treated with one or more applications of the hypersensitive response elicitor in accordance with the present invention to impart stress resistance. Such plants may also be treated with conventional plant treatment agents (e.g., insecticides, fertilizers, etc.).

## EXAMPLES

### **Example 1 - Hypersensitive Response Elicitor-Treated Cotton is More Resistant to the Damage Caused by Insecticide Stress**

Aphids (*Aphis gossypii*) infect cotton during the entire growth season. The damage of aphid infection ranges from honeydew deposit that contaminates the lint and reduces crop value to defoliation that reduces or destroys crops. To protect plants from aphid infection, cotton is usually sprayed with insecticides, for example Asana XL when the infection pressure is not very high, and Admire when the infestation pressure is high. The effect of a hypersensitive response elicitor on aphids in cotton was studied by a trial involving a randomized complete block design. This

- 42 -

involved treatment with *Erwinia amylovora* hypersensitive response elicitor (i.e. HP-1000™) at 20, 60, and 80 ppm and a chemical insecticide, Asana XL, at 8 oz./ac. Each treatment involved foliar application beginning at cotyledon to three true leaves and thereafter at 14 day intervals using a backpack sprayer. Aphid counts and overall growth of the cotton were made immediately prior to spray application at 14, 28, 35, and 42 days after the first treatment ("DAT 1"). Twenty-five randomly selected leaves per plot were collected at the first three sampling dates and the leaves per plot at the final sampling date.

## 10 Results

1. Aphid control: The number of aphids in the hypersensitive response elicitor-treated cotton were significantly reduced in comparison to the chemical treated cotton (see Table 1).

15 Table 1. Aphid count per leaf on cotton after treatment with Asana XL® or HP-1000™

Treatment	Rate <sup>2</sup>	Number of aphids per leaf <sup>1</sup>			
		No. sprays applied/days after treatment			
		1/14DAT1	2/28DAT1	3/35DAT1	4/42DAT1
Asana XL®	8 oz/ac	0.2 a	32.2 a	110.0 a	546.9 a
HP-1000™	20 µg/ml	0.2 a	7.8 b	22.9 b	322.1 a
HP-1000™	60 µg/ml	0.1 a	4.9 b	34.6 b	168.3 a
HP-1000™	80 µg/ml	0.0 a	2.7 b	25.8 b	510.2 a

<sup>1</sup>Means followed by different letters are significantly different according to Duncan's MRT, P=0.05. <sup>2</sup>Rate for Asana XL® is for formulated product, rate for HP-1000™ is for active ingredient (a.i.).

At 14 days after DAT 1, aphid counts were relatively low across all of the treatments, but by 28 days after DAT 1 (by which time two sprayings had been applied), the number of aphids per leaf were significantly greater in Asana XL-treated plants compared to the hypersensitive response elicitor-treated cottons. By 35 days after DAT 1 (by which time three sprayings had been applied), aphid counts had risen for all treatments, yet aphid counts per leaf were still significantly lower for hypersensitive response elicitor-treated cotton compared to the Asana XL treatment. Finally, at 42 days after DAT 1 (by which time four sprayings had been applied), the number of aphids per leaf had increased to a level that threatened to overwhelm the

plants even when treated with the standard chemical insecticide. To save the trial, another chemical, Pravado (Admire), was applied to all plots to eradicate aphids from the field.

2. Hypersensitive response elicitor-treated cotton was more resistant to the damage caused by Pravado (Admire) and Asana. After the second chemical spraying, it was observed that cotton plants were stress shocked by the insecticides. The cotton plants previously treated with Asana and untreated control were defoliated. On most of the chemical-treated cotton, there were no leaves, or very few leaves, in the lower portion of plants. However, the hypersensitive response elicitor-treated plants, especially the plot where hypersensitive response elicitor was applied at 80 ppm, had no defoliation and the cotton plants were vigorous and healthy. By counting the number of mature balls, it clearly showed that hypersensitive response elicitor-treated plants (at 80 ppm) had more ball setting than chemical and untreated control (Table 2), indicating that hypersensitive response elicitor-treated plants were more tolerant to the stress caused by insecticide.

Table 2. Number of Formed Cotton Balls Counted on Ten Plants in Each of Four Replicates Per Treatment.

<u>Treatment</u>	<u>No. balls/10 plants/replicate</u>
UTC	28
Chemical standard	6
Hypersensitive Response Elicitor	35

**Example 2 - Hypersensitive Response Elicitor-Treated Cucumbers are More Resistant to Drought**

A cucumber field trial was set up to test the effect of *Erwinia amylovora* hypersensitive response elicitor on disease control, tolerance to drought stress, and yield. Three different rates were tested, there at 15, 30, and 60 µg/ml. In addition to hypersensitive response elicitor treatment, there was an untreated control. Each treatment contained three replicate plots. When the first true leaf emerges, hypersensitive response elicitor was sprayed with a back bag sprayer. The second spray was applied ten days after the first spray. The third application was right after

the recovery of cucumber seedlings after the transplanting to the field. Individual treatment was randomly assigned in the field.

When the first true leaf emerged (Day 0), a first application was sprayed. Usually cucumber seedlings are transplanted when seedlings show two true leaves. It has been known that the recovery rate after the transplanting is closely related to the size of the seedlings. Because of the drought, the seedlings were maintained in the nursery for an extra ten days and the second spray was applied on Day 10. Two days after the second spray, the plants were transplanted into fields and covered with plastic sheets. The plants had 4 – 5 true leaves.

10

### Result

The recovery rate of the transplanted cucumber seedlings was higher for the hypersensitive response elicitor-treated plants than for the untreated control. More than 80% of the hypersensitive response elicitor-treated cucumber seedlings survived, while only 57% untreated plants survived.

15

Throughout the growth season, there was a serious drought problem. Early field visits indicated that hypersensitive response elicitor-treated plants had more root mass and better over-all growth. Hypersensitive response elicitor-treated cucumber started to flower 14 days earlier than untreated control cucumber. The early flowering resulted in an earlier harvest. In the first harvest, more than 0.4 kilograms of cucumber fruits per plant were harvested from the hypersensitive response elicitor-treated cucumbers; however, virtually no fruit was harvested from untreated control. By the end of the season, untreated plants died due to severe drought, but hypersensitive response elicitor-treated plants were still alive and had one more harvest.

20

25

The final yield was significantly different between hypersensitive response elicitor-treated and untreated plants. Hypersensitive response elicitor administered at the rate of 30 ppm produced three times greater yield than the control plants (Table 3).

30

Table 3. Yield Increase of Cucumber Fruit from Hypersensitive Response Elicitor Treated Plants

Treatment	Replicate	kg/plant	Yield/Replicate	% of the Yield Increase
HP 15	I	1.25	37.5	
	II	1.00	30.0	103.8
	III	1.21	36.3	241
HP 30	I	1.54	46.2	
	II	1.43	42.9	133.2
	III	1.47	44.1	339
Control	I	0.43	12.9	
	II	0.41	12.3	39.3
	III	0.47	14.1	

5

The increased yield was partially attributed to hypersensitive response elicitor-induced growth enhancement and partially resulted from more tolerance of hypersensitive response elicitor-treated cucumber to drought, because usually the yield increase from hypersensitive response elicitor-induced growth enhancement is between 10-40%.

10

**Example 3 - Hypersensitive Response Elicitor-Treated Pepper is More Tolerant to Herbicide Stress**

15

Pepper seedlings were drenched with hypersensitive response elicitor at 20 ppm seven days before transplanting, sprayed seven days after the transplanting, and then, sprayed every fourteen days. Standard chemicals, Brave, Maneb, Kocide, and Admire, were used for the rest of the treatment. In addition to early growth enhancement, which resulted in a higher yield, larger fruit, and resistance to several diseases, hypersensitive response elicitor-treated pepper was more tolerant to herbicide damage. The pepper field was applied with the herbicide SENCOR which is not labeled for pepper. This herbicide is known to cause severe foliar damage to pepper in chemically-treated plants but not with hypersensitive response elicitor-treated plants.

20

25

The difference between the adverse effect of the herbicide on the hypersensitive response elicitor and non-hypersensitive response elicitor treated plants is dramatic. See Table 4 below. Thirty-nine of the 60 elicitor-treated plants showed only minor damage by the herbicide, the damaged leaves were less than 20%. In

contrast, 53 out of the 60 chemically-treated pepper plants had severe damage, 40-57% of the leaves were damaged, and 20 plants were dead. The ability of hypersensitive response elicitors to help crops withstand the phytotoxic effects of a herbicide is very important benefit to in agricultural industry.

5

Table 4. Hypersensitive Response Elicitor-Treated Peppers are More Tolerant to Herbicide Damage.

Treatment	Damage Rating						Damage Index %
	1	2	3	4	5	6	
Hypersensitive Response Elicitor	1	38	17	3	1	0	41
Chemicals	0	1	6	16	19	18	87

15

Damage Rating: 1. No damage; 2. 0-20% leaves damaged; 3. 20-40% leaves damaged; 4. 40-50% leaves damaged; 6. More than 75% leaves damaged or entire plant dead.

20

Damage index = sum of each rating times the number of plants under the rating scale, divided by total number of plants times 6.

Damage index for hypersensitive response elicitor-treated plants =  $\frac{1 \times 1 + 2 \times 38 + 3 \times 17 + 4 \times 3 + 5 \times 1 + 6 \times 0}{6 \times 60} \times 100\% = 41\%$

25

#### **Example 4 - Hypersensitive Response Elicitor-Treated Pepper is More Tolerant to Herbicide Stress under Controlled Experimental Conditions**

30

A field trial was conducted to test if hypersensitive elicitor treated pepper would be more tolerant to herbicide stress. The trial contains 6 treatments and 4 replicates for each treatment. The treatments are described as follows:

1. Control, the peppers were neither treated by a hypersensitive response ("HR") elicitor nor by LEXONE™ herbicide (DuPont Agricultural Products, Wilmington, Delaware).
2. Control pepper with application of 0.15 pound LEXONE™ herbicide /acre.
3. Control pepper with application of 0.3 pound LEXONE™ herbicide /acre.

40



- 47 -

4. HR elicitor treatment with no application of LEXONE™ herbicide using a formulated product known as MESSENGER™ biopesticide (Eden Bioscience Corporation, Bothell, Washington) containing 3% HR elicitor protein was used.

5 5. HR elicitor treatment with application of 0.15 pound LEXONE™ herbicide /acre.

6. HR elicitor treatment with application of 0.3 pound LEXONE™ herbicide /acre.

LEXONE™ contains the same active ingredient as SENCOR™ herbicide (Bayer, Kansas City, Missouri) used in Example 3. Pepper seedlings were drenched with MESSENGER™ solution at the concentration of HR elicitor protein of about 20 ppm seven days before transplanting into the field and then sprayed every 14 days after the transplanting. LEXONE was applied at high (0.3 pound/acre) and low levels (0.15 pound/acre). 50 gallon water and 100 mL of the herbicide solution was introduced into the root zone of each plant in the respective treatment five weeks after transplant into the field.

The treatments were evaluated for the percent of chlorosis caused by the LEXONE™ herbicide application and for the pepper yield. HR elicitor-treated plants exposed to the high rate of herbicide had significantly less chlorosis and produced 108 % more fruit in comparison to the non-hypersensitive response elicitor treated plants exposed to the same amount of herbicide. See Tables 5 and 6 below. There was no significant difference in the reduction of chlorosis at the low rate of herbicide between the HR elicitor treated and non-HR elicitor treated peppers. However, the HR elicitor treated plants produced 15% more fruit than the corresponding control plants exposed to the same amount of herbicide. There was no chlorosis in either the check or HR elicitor-treated plants that did not receive LEXONE™ herbicide treatment.

The HR elicitor treated plants were much less severely affected by the herbicide application than the respective control plants at the high rate of herbicide. However, the amount of visual chlorosis was similar at the low rate for both the check and HR elicitor-treated plants. More importantly, the yields from both the high and low rate herbicide treatments of HR elicitor treated plants were less severely effected

by the herbicide than the checks. These findings further confirm that HR elicitors can help crops withstand the phytotoxic effects of herbicides and are very beneficial to the agricultural industry.

5 Table 5. Reduction of Foliar Chlorosis and Increase in Yield in Hypersensitive Response Elicitor Treated Plants after Exposure to LEXONE™ Herbicide

Treatment	Percent foliar chlorosis and yield of pepper						% difference from the respective control
	A	B	C	D	E	Yield (pound)	
6 (MESSENGER™ + High rate LEXONE™)	13.75	30.00	37.50	36.25	40.00	8.31	108 %
3 (High rate LEXONE™)	26.25	43.75	51.25	50.00	51.25	4.00	-
5 (MESSENGER™ + low rate LENOXETM)	16.25	22.50	28.75	23.75	27.50	8.00	15 %
2 (LENOXETM)	12.50	20.00	25.00	25.00	23.75	6.81	-

10 Table 6. Weight of Harvested Peppers Increased in Hypersensitive Response Elicitor Treated Plants after Exposure to LEXONE™ Herbicide Compared to Check Plants.

Treatment	Weight of peppers harvested 12/1/98 in pounds
HP20 + high rate LEXONE™	8.31
Check + high rate LEXONE™	4.00
HP20 + low rate LEXONE™	8.00
Check + low rate LEXONE™	6.81

15 **Example 5 - Hypersensitive Response Elicitor-Treated Cotton is More Tolerant to Drought Stress**

A non-irrigated cotton trial experienced 26 consecutive days of drought. The average daily heat index was near or over 100 degrees F, adding to the stress placed on the plants in the field.

20

Observations in the field indicated that plants treated with HR elicitor at the concentration of 15 ppm (2.2 oz formulated product, MESSENGER™ containing 3 % active ingredient HR elicitor protein) were more vigorous and had less defoliation than the check plants as a result of the heat and drought stress. Equal numbers of plants from the MESSENGER™-treated and the non-MESSENGER™ treated plots were carefully removed from the field and mapped for the number of nodes and bolls by position. The plants were also weighed on a Metler analytical scale to determine whole plant, root and shoot weights.

MESSENGER™ treated plants survived the heat and drought stresses much better than the untreated plants did. Plants treated with MESSENGER™ had 37.6% more root and shoot mass than the check plants (Table 7). The MESSENGER™ treated plants also had significantly more cotton bolls than the check plants (Table 8). The number of cotton bolls from positions 1 and 2 have a significant contribution to the overall yield. Table 8 showed that MESSENGER™ treated plants had 47% more bolls in positions 1 and 2 and 57% more boll from a whole plant in comparison to the yield achieved using a grower standard treatment (i.e. with no MESSENGER™ treatment). A common reaction to stress in cotton is for the plant to abort bolls. The results indicate that MESSENGER™-treated plants are more tolerant to the drought stress.

Table 7. Weight per Plant of Non-Irrigated Cotton Following 26 Days of Drought.

Treatment	Root weight (pond/plant)	%Difference	Shoot weight (pond/plant)	% difference	Whole plant weight (pond/plant)	% difference
MESSENGER™ 2.2 oz/acre	0.041 a*	37.6 %	0.505 a	37.5 %	0.546	37.5 %
Control (Grower standard)	0.0298 b		0.367 b		0.397	
Level of statistically significant	P=0.119		P=0.034			P=0.033

\* Same letter indicates no statistical difference between the two treatments at the defined level; different letter indicates a statistical difference between the two treatments at the defined level.

Table 8. Number of Bolls per 5 Plants at the Number 1 & 2 positions, and Total Number of Bolls from Whole Plants in Non-irrigated Cotton Following 26 days of drought.

5

Treatment	Avg. # bolls in the #1 & 2 position	Percent difference	Avg. # of total bolls per 5 plant	Percent difference
MESSENGER™ 2.2 OZ.	18.4 a	+46.0%	21.4 a	+57.0%
Check	12.6 b		13.6 b	-
Statistically significant level	P=0.032		P=0.01	

\* Same letter indicates no statistical difference between the two treatments at the defined level; different letter indicates a statistical difference between the two treatments at the defined level.

10 **Example 6 - Hypersensitive Response Elicitor-Treated Tomato is More Tolerant to Calcium Deficiency**

Calcium is an important element for plant physiology and development. A deficiency in calcium can cause several plant diseases. For example, blossom-end rot is caused by a localized calcium deficiency in the distal end of the tomato fruit. Because calcium is not a highly mobile element, a deficiency can occur with a fluctuation in water supply. In the past, tomato growers experienced higher level of blossom-end rot during dry weather conditions when infrequent rains storms dumped a lot of water and then return to a hot and dry condition quickly. Lowering or raising the irrigation water table erratically during a dry and hot growing season can also increase the disease.

A field trial was designed to test if HR elicitor protein-treated tomato can be more tolerant to the calcium deficiency under a dry hot growing season. MESSENGER™, the formulated product containing 3% HR elicitor, was used for the trial. The application rate of the MESSENGER™ was 2.27 oz per care. The first spray of MESSENGER™ was carried out 7 days before the transplanting and then every 14-days after transplanting. MESSENGER™-treated tomatoes were compared with a standard grower treatment not utilizing MESSENGER™. Each treatment had 4 replicates.

30 The number of infected fruit was counted from a 100 square foot field. The rot typically begins with light tan water soaked lesions, which then enlarge, and then turn black. In a survey, about 20% of the fruits were infected. Severe end-rot

- 51 -

symptoms occurred in the standard treatment; however, an average of only 2.5 % of the fruit was infected in the MESSENGER™-treated plants. The harvest data showed that MESSENGER™-treated plants had 8% more marketable fruit (Table 9). The test results demonstrated that MESSENGER™-treatment can reduce the stress resulting from calcium deficiency and increase plant resistance to blossom-end rot.

**Table 9. Hypersensitive Response Elicitor Treatment Reduced Blossom-End Rot Infection, Increased Yield of Tomato Fruit**

Treatment	Blossom-End Infected Fruit*				Tomato Fruit Yield	
	Rep I	Rep II	Rep III	Rep IV	Bin/Acre	% Difference
MESSENGER™	0	9	0	1	35	8
Standard Treatment)	24	22	16	17	31.5	-

\*The data were collected from the fruits in 100 square foot plot

Although the invention has been described in detail for the purpose of illustration, it is understood that such detail is solely for that purpose, and variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention which is defined by the following claims.

**WHAT IS CLAIMED:**

1. A method of imparting stress resistance to plants comprising:  
applying a hypersensitive response elicitor protein or  
5 polypeptide in a non-infectious form to a plant or plant seed under conditions  
effective to impart stress resistance.
2. A method according to claim 1, wherein the stress resistance is  
resistance to a stress selected from the group consisting of climated related stress, air  
10 pollution stress, chemical stress, and nutritional stress.
3. A method according to claim 2, wherein the stress is chemical  
stress where the chemical is selected from the group consisting of insecticides,  
fungicides, herbicides, and heavy metals.
- 15 4. A method according to claim 2, wherein the stress is climate-  
related stress selected from the group consisting of drought, water, frost, cold  
temperature, high temperature, excessive light, and insufficient light.
- 20 5. A method according to claim 2, wherein the stress is air  
pollution stress selected from the group consisting of carbon dioxide, carbon  
monoxide, sulfur dioxide, NO<sub>x</sub>, hydrocarbons, ozone, ultraviolet radiation, and acidic  
rain.
- 25 6. A method according to claim 2, wherein the stress is nutritional  
stress where the nutritional stress is caused by fertilizer, micronutrients, or  
macronutrients.
- 30 7. A method according to claim 1, wherein the hypersensitive  
response elicitor protein or polypeptide is derived from *Erwinia*, *Pseudomonas*,  
*Xanthamonas*, *Phythophthera*, or *Clavibacter*.

8. A method according to claim 7, wherein the hypersensitive response elicitor protein or polypeptide is derived from *Erwinia amylovora*, *Erwinia carotovora*, *Erwinia chrysanthemi*, and *Erwinia stewartii*.
- 5 9. A method according to claim 7, wherein the hypersensitive response elicitor protein or polypeptide is derived from *Pseudomonas syringae* or *Pseudomonas solanacearum*.
- 10 10. A method according to claim 7, wherein the hypersensitive response elicitor protein or polypeptide is derived from a *Xanthomonas* species.
11. A method according to claim 7, wherein the hypersensitive response elicitor protein or polypeptide is derived from a *Phytophthora*.
- 15 12. A method according to claim 7, wherein the hypersensitive response elicitor protein or polypeptide is derived from *Clavibacter michiganensis* subsp. *sepedonicus*.
- 20 13. A method according to claim 1, wherein plants are treated during said applying.
14. A method according to claim 1, wherein plant seeds are treated during said applying, said method further comprising:  
planting the seeds treated with the hypersensitive response  
25 elicitor protein or polypeptide in natural or artificial soil and propagating plants from seeds planted in soil.
15. A method according to claim 1, wherein the plant is selected from the group consisting of alfalfa, rice, wheat, barley, rye, cotton, sunflower,  
30 peanut, corn, potato, sweet potato, bean pea, chicory, lettuce, endive, cabbage, brussel sprout, beet, parsnip, cauliflower, broccoli, turnip, radish, spinach, onion, garlic, eggplant, pepper, celery, carrot, squash, pumpkin, zucchini, cucumber, apple, pear,

- 54 -

melon, citrus, strawberry, grape, raspberry, pineapple, soybean, tobacco, tomato, sorghum, and sugarcane.

16. A method according to claim 1, wherein the plant is selected  
5 from the group consisting of *Arabidopsis thaliana*, *Saintpaulia*, petunia, pelargonium, poinsettia, chrysanthemum, carnation, and zinnia.

17. A method of imparting stress resistance to plants comprising:  
providing a transgenic plant or plant seed transformed with a  
10 DNA molecule which encodes for a hypersensitive response elicitor protein or polypeptide and  
growing the transgenic plant or plants produced from the  
transgenic plant seeds under conditions effective to impart stress resistance.

18. A method according to claim 17, wherein a transgenic plant is  
15 provided.

19. A method according to claim 17, wherein a transgenic plant  
seed is provided, said method further comprising:  
20 planting the transgenic seeds in natural or artificial soil and  
propagating plants from seeds planted in soil..

20. A method according to claim 17, wherein the stress resistance  
is resistance to a stress selected from the group consisting of climated related stress,  
25 air pollution stress, chemical stress, and nutritional stress.

21. A method according to claim 20, wherein the stress is chemical  
stress where the chemical is selected from the group consisting of insecticides,  
fungicides, herbicides, and heavy metals.

30



22. A method according to claim 20, wherein the stress is climate-related stress selected from the group consisting of drought, water, frost, cold temperature, high temperature, excessive light, and insufficient light.
- 5 23. A method according to claim 20, wherein the stress is air pollution stress selected from the group consisting of carbon dioxide, carbon monoxide, sulfur dioxide, NO<sub>x</sub>, hydrocarbons, ozone, ultraviolet radiation, and acidic rain.
- 10 24. A method according to claim 20, wherein the stress is nutritional stress where the nutritional stress is caused by fertilizer, micronutrients, or macronutrients.
- 15 25. A method according to claim 20, wherein the hypersensitive response elicitor protein or polypeptide is derived from *Erwinia*, *Pseudomonas*, *Xanthomonas*, *Phytophthora*, or *Clavibacter*.
- 20 26. A method according to claim 25, wherein the hypersensitive response elicitor protein or polypeptide is derived from *Erwinia amylovora*, *Erwinia carotovora*, *Erwinia chrysanthemi*, and *Erwinia stewartii*.
- 25 27. A method according to claim 25, wherein the hypersensitive response elicitor protein or polypeptide is derived from *Pseudomonas syringae* or *Pseudomonas solanacearum*.
28. A method according to claim 25, wherein the hypersensitive response elicitor protein or polypeptide is derived from a *Xanthomonas* species.
- 30 29. A method according to claim 20, wherein the plant is selected from the group consisting of alfalfa, rice, wheat, barley, rye, cotton, sunflower, peanut, corn, potato, sweet potato, bean pea, chicory, lettuce, endive, cabbage, brussel sprout, beet, parsnip, cauliflower, broccoli, turnip, radish, spinach, onion, garlic,

- 56 -

eggplant, pepper, celery, carrot, squash, pumpkin, zucchini, cucumber, apple, pear, melon, citrus, strawberry, grape, raspberry, pineapple, soybean, tobacco, tomato, sorghum, and sugarcane.

- 5                    30.    A method according to claim 20, wherein the plant is selected from the group consisting of *Arabidopsis thaliana*, *Saintpaulia*, petunia, pelargonium, poinsettia, chrysanthemum, carnation, and zinnia.

## SEQUENCE LISTING

&lt;110&gt; Eden Bioscience Corporation

<120> HYPERSENSITIVE RESPONSE ELICITOR-INDUCED STRESS  
RESISTANCE

&lt;130&gt; 21829/42

&lt;140&gt;

&lt;141&gt;

&lt;150&gt; 60/107,243

&lt;151&gt; 1998-11-05

&lt;160&gt; 18

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 338

&lt;212&gt; PRT

&lt;213&gt; Erwinia chrysanthemi

&lt;400&gt; 1

Met	Gln	Ile	Thr	Ile	Lys	Ala	His	Ile	Gly	Gly	Asp	Leu	Gly	Val	Ser
1				5					10					15	

Gly	Leu	Gly	Ala	Gln	Gly	Leu	Lys	Gly	Leu	Asn	Ser	Ala	Ala	Ser	Ser
		20						25					30		

Leu	Gly	Ser	Ser	Val	Asp	Lys	Leu	Ser	Ser	Thr	Ile	Asp	Lys	Leu	Thr
		35					40					45			

Ser	Ala	Leu	Thr	Ser	Met	Met	Phe	Gly	Gly	Ala	Leu	Ala	Gln	Gly	Leu
	50					55					60				

Gly	Ala	Ser	Ser	Lys	Gly	Leu	Gly	Met	Ser	Asn	Gln	Leu	Gly	Gln	Ser
65					70					75				80	

Phe	Gly	Asn	Gly	Ala	Gln	Gly	Ala	Ser	Asn	Leu	Leu	Ser	Val	Pro	Lys
				85					90					95	

Ser	Gly	Gly	Asp	Ala	Leu	Ser	Lys	Met	Phe	Asp	Lys	Ala	Leu	Asp	Asp
			100					105					110		

Leu	Leu	Gly	His	Asp	Thr	Val	Thr	Lys	Leu	Thr	Asn	Gln	Ser	Asn	Gln
		115					120					125			



Leu Ala Asn Ser Met Leu Asn Ala Ser Gln Met Thr Gln Gly Asn Met  
 130 135 140  
 Asn Ala Phe Gly Ser Gly Val Asn Asn Ala Leu Ser Ser Ile Leu Gly  
 145 150 155 160  
 Asn Gly Leu Gly Gln Ser Met Ser Gly Phe Ser Gln Pro Ser Leu Gly  
 165 170 175  
 Ala Gly Gly Leu Gln Gly Leu Ser Gly Ala Gly Ala Phe Asn Gln Leu  
 180 185 190  
 Gly Asn Ala Ile Gly Met Gly Val Gly Gln Asn Ala Ala Leu Ser Ala  
 195 200 205  
 Leu Ser Asn Val Ser Thr His Val Asp Gly Asn Asn Arg His Phe Val  
 210 215 220  
 Asp Lys Glu Asp Arg Gly Met Ala Lys Glu Ile Gly Gln Phe Met Asp  
 225 230 235 240  
 Gln Tyr Pro Glu Ile Phe Gly Lys Pro Glu Tyr Gln Lys Asp Gly Trp  
 245 250 255  
 Ser Ser Pro Lys Thr Asp Asp Lys Ser Trp Ala Lys Ala Leu Ser Lys  
 260 265 270  
 Pro Asp Asp Asp Gly Met Thr Gly Ala Ser Met Asp Lys Phe Arg Gln  
 275 280 285  
 Ala Met Gly Met Ile Lys Ser Ala Val Ala Gly Asp Thr Gly Asn Thr  
 290 295 300  
 Asn Leu Asn Leu Arg Gly Ala Gly Gly Ala Ser Leu Gly Ile Asp Ala  
 305 310 315 320  
 Ala Val Val Gly Asp Lys Ile Ala Asn Met Ser Leu Gly Lys Leu Ala  
 325 330 335  
 Asn Ala

&lt;210&gt; 2

&lt;211&gt; 2141

&lt;212&gt; DNA

&lt;213&gt; Erwinia chrysanthemi



&lt;400&gt; 2

```

cgattttacc cgggtgaacg tgctatgacc gacagcatca cgggtattcga caccgttacg 60
gcgtttatgg ccgcgatgaa ccggcatcag gcggcgcgct ggtcgccgca atccggcgctc 120
gatctgggat ttcagtttgg ggacaccggg cgtgaactca tgatgcagat tcagccgggg 180
cagcaatata ccggcatggt gcgcacgctg ctgcgtcgtc gttatcagca ggccggcagag 240
tgcgatggct gccatctgtg cctgaacggc agcgatgtat tgatcctctg gtggccgctg 300
ccgtcgggac ccggcagtta tccgcagggt atcgaacggt tgtttgaact ggccgggaatg 360
acgttgccgt cgctatccat agcaccgacg gcgcgtccgc agacagggaa cggacgcgcc 420
cgatcattaa gataaaggcg gcttttttta ttgcaaaacg gtaacgggtg ggaaccgttt 480
caccgtcggc gtcactcagt aacaagtata catcatgatg cctacatcgg gatcggcgctg 540
ggcatccgtt gcagataact ttgcgaacac ctgacatgaa tgaggaaacg aaattatgca 600
aattacgatc aaagcgcaca tccggcggtg tttggcgctc tccgggtctg ggctgggtgc 660
tcagggactg aaaggactga attccgcggc ttcacgctg gggtccagcg tggataaact 720
gagcagcacc atcgataagt tgacctccgc gctgacttcg atgatgtttg gcggcgcgct 780
ggcgcgaggg ctggcgccca gctcgaagg gctggggatg agcaatcaac tgggccagtc 840
tttcggcaat ggcgcgcgag gtgcgagcaa cctgctatcc gtaccgaaat ccggcgggcg 900
tgcgttgtca aaaatgtttg ataaagcgct ggacgatctg ctgggtcatg acaccgtgac 960
caagctgact aaccagagca accaactggc taattcaatg ctgaacgcca gccagatgac 1020
ccagggtaat atgaatgcgt tccgcagcgg tgtgaacaac gcaactgtcg ccattctcgg 1080
caacggtctc ggccagtcga tgagtggctt ctctcagcct tctctggggg caggcggtt 1140
gcagggcctg agcggcgcgg gtgcattcaa ccagttgggt aatgccatcg gcatggcgct 1200
ggggcagaat gctgcgctga gtgcgttgag taacgtcagc acccacgtag acggtataca 1260
ccgccacttt gtagataaag aagatcgcg catggcgaaa gagatcggcc agtttatgga 1320
tcagtatccg gaaatattcg gtaaaccgga ataccagaaa gatggctgga gttcgccgaa 1380
gacggacgac aaatcctggg ctaaagcgct gagtaaacgg gatgatgacg gtatgaccgg 1440
cgccagcatg gacaaattcc gtcaggcgat gggtatgatc aaaagcgcg tggcggtgga 1500
taccggcaat accaacctga acctgcgtgg cgccggcggt gcacgctgg gtatcgatgc 1560
ggctgtcgtc ggcgataaaa tagccaacat gtcgctgggt aagctggcca acgcctgata 1620
atctgtgctg gcctgataaa gcggaaacga aaaaagagac ggggaagcct gtctcttttc 1680
ttattatgcg gtttatgcgg ttacctggac cggttaatca tcgtcatcga tctggtacaa 1740
acgcacattt tcccgttcac tcgcgtcggt acgcgccaca atcgcgatgg catcttcctc 1800
gtcgctcaga ttgcgcggt gatggggaac gccgggtgga atatatagaa actcgccggc 1860
cagatggaga cacgtctgcg ataaatctgt gccgtaacgt gtttctatcc gcccttttag 1920
cagatagatt gcggtttcgt aatcaacatg gtaatgcggt tccgcctgtg cgccggccgg 1980
gatcaccaca atattcatag aaagctgtct tgcacctacc gtatcgcggg agataccgac 2040
aaaataggcg agtttttgcg tggtatccgt ggggtgttcc ggcctgacaa tcttgagttg 2100
gttcgtcatc atctttctcc atctggcgga cctgatcggt t 2141

```

&lt;210&gt; 3

&lt;211&gt; 403

&lt;212&gt; PRT

<213> *Erwinia amylovora*

&lt;400&gt; 3

```

Met Ser Leu Asn Thr Ser Gly Leu Gly Ala Ser Thr Met Gln Ile Ser
1              5              10              15

```





11 Gly Gly Ala Gly Gly Asn Asn Gly Leu Leu Gly Thr Ser Arg Gln  
                     20                    25                    30  
  
 Asn Ala Gly Leu Gly Gly Asn Ser Ala Leu Gly Leu Gly Gly Gly Asn  
                     35                    40                    45  
  
 Gln Asn Asp Thr Val Asn Gln Leu Ala Gly Leu Leu Thr Gly Met Met  
                     50                    55                    60  
  
 Met Met Met Ser Met Met Gly Gly Gly Gly Leu Met Gly Gly Gly Leu  
                     65                    70                    75                    80  
  
 Gly Gly Gly Leu Gly Asn Gly Leu Gly Gly Ser Gly Gly Leu Gly Glu  
                     85                    90                    95  
  
 Gly Leu Ser Asn Ala Leu Asn Asp Met Leu Gly Gly Ser Leu Asn Thr  
                     100                    105                    110  
  
 Leu Gly Ser Lys Gly Gly Asn Asn Thr Thr Ser Thr Thr Asn Ser Pro  
                     115                    120                    125  
  
 Leu Asp Gln Ala Leu Gly Ile Asn Ser Thr Ser Gln Asn Asp Asp Ser  
                     130                    135                    140  
  
 Thr Ser Gly Thr Asp Ser Thr Ser Asp Ser Ser Asp Pro Met Gln Gln  
                     145                    150                    155                    160  
  
 Leu Leu Lys Met Phe Ser Glu Ile Met Gln Ser Leu Phe Gly Asp Gly  
                     165                    170                    175  
  
 Gln Asp Gly Thr Gln Gly Ser Ser Ser Gly Gly Lys Gln Pro Thr Glu  
                     180                    185                    190  
  
 Gly Glu Gln Asn Ala Tyr Lys Lys Gly Val Thr Asp Ala Leu Ser Gly  
                     195                    200                    205

---

Leu Met Gly Asn Gly Leu Ser Gln Leu Leu Gly Asn Gly Gly Leu Gly  
                     210                    215                    220  
  
 Gly Gly Gln Gly Gly Asn Ala Gly Thr Gly Leu Asp Gly Ser Ser Leu  
                     225                    230                    235                    240  
  
 Gly Gly Lys Gly Leu Gln Asn Leu Ser Gly Pro Val Asp Tyr Gln Gln  
                     245                    250                    255  
  
 Leu Gly Asn Ala Val Gly Thr Gly Ile Gly Met Lys Ala Gly Ile Gln  
                     260                    265                    270



Ala Leu Asn Asp Ile Gly Thr His Arg His Ser S r Thr Arg Ser Ph  
 275 280 285

Val Asn Lys Gly Asp Arg Ala Met Ala Lys Glu Ile Gly Gln Phe Met  
 290 295 300

Asp Gln Tyr Pro Glu Val Phe Gly Lys Pro Gln Tyr Gln Lys Gly Pro  
 305 310 315 320

Gly Gln Glu Val Lys Thr Asp Asp Lys Ser Trp Ala Lys Ala Leu Ser  
 325 330 335

Lys Pro Asp Asp Asp Gly Met Thr Pro Ala Ser Met Glu Gln Phe Asn  
 340 345 350

Lys Ala Lys Gly Met Ile Lys Arg Pro Met Ala Gly Asp Thr Gly Asn  
 355 360 365

Gly Asn Leu Gln Ala Arg Gly Ala Gly Gly Ser Ser Leu Gly Ile Asp  
 370 375 380

Ala Met Met Ala Gly Asp Ala Ile Asn Asn Met Ala Leu Gly Lys Leu  
 385 390 395 400

Gly Ala Ala

<210> 4

<211> 1288

<212> DNA

<213> *Erwinia amylovora*

<400> 4

aagcttcggc atggcacgtt tgaccgttgg gtcggcaggg tacgtttgaa ttattcataa 60

---

gaggaatacg ttatgagtct gaatacaagt gggctgggag cgtcaacgat gcaaatttct 120

atcggcgggtg cgggcggaaa taacgggttg ctgggtacca gtcgccagaa tgctgggttg 180

ggtggcaatt ctgcaactggg gctgggcggc ggtaatcaaa atgataccgt caatcagctg 240

gctggcttac tcaccggcat gatgatgatg atgagcatga tgggcgggtg tgggctgatg 300

ggcgggtggct taggcgggtg cttaggtaat ggcttgggtg gctcaggtgg cctgggcgaa 360

ggactgtcga acgcgctgaa cgatatgtta ggcggttcgc tgaacacgct gggctcgaaa 420

ggcggcaaca ataccacttc aacaacaaat tccccgctgg accaggcgct ggggtattaac 480

tcaacgtccc aaaacgacga ttccacctcc ggcacagatt ccacctcaga ctccagcgac 540

ccgatgcagc agctgctgaa gatgttcagc gagataatgc aaagcctgtt tgggtgatggg 600

caagatggca cccagggcag ttctctctggg ggcaagcagc cgaccgaagg cgagcagaac 660

gcctataaaa aaggagtcac tgatgcgctg tcgggcctga tgggtaatgg tctgagccag 720

ctccttggca acgggggact gggaggtggg cagggcggtg atgctggcac gggctcttgac 780

ggttcgtcgc tgggcggcaa agggctgcaa aacctgagcg ggccgggtgga ctaccagcag 840



```

ttaggtaacg ccggtgggtac cgggtatcgggt atgaaagcgg gcattcaggg gctgaatgat 900
atcgggtacgc acaggcacag ttcaaccggt tctttcgtca ataaaggcga tcgggcgatg 960
gcgaaggaaa tcgggtcagtt catggaccag tatcctgagg tgtttggcaa gccgcagtac 1020
cagaaaggcc cgggtcagga ggtgaaaacc gatgacaaat catgggcaaa agcactgagc 1080
aagccagatg acgacggaat gacaccagcc agtatggagc agttcaacaa agccaagggc 1140
atgatcaaaa ggcccatggc ggggtgatacc ggcaacggca acctgcaggg acgcggtgcc 1200
ggtgggttctt cgctgggtat tgatgccatg atggccgggtg atgccattaa caatatggca 1260
cttggcaagc tgggcgcggc ttaagctt 1288

```

&lt;210&gt; 5

&lt;211&gt; 1344

&lt;212&gt; DNA

<213> *Erwinia amylovora*

&lt;400&gt; 5

```

atgtcaattc ttacgcttaa caacaatacc tcgtcctcgc cgggtctggt ccagtcgggg 60
ggggacaacg ggcttggtgg tcataatgca aattctgcgt tggggcaaca acccatcgat 120
cggcaaacca ttgagcaaat ggctcaatta ttggcggaac tggttaaagtc actgctatcg 180
ccacaatcag gtaatgcggc aaccggagcc ggtggcaatg accagactac aggagttggt 240
aacgctggcg gcctgaacgg acgaaaaggc acagcaggaa cactccgca gtctgacagt 300
cagaacatgc tgagtgaatg gggcaacaac gggctggatc aggccatcac gcccgatggc 360
cagggcgggc ggcagatcgg cgataatcct ttactgaaag ccatgctgaa gcttattgca 420
cgcatgatgg acggccaaag cgatcagttt ggccaacctg gtacgggcaa caacagtgcc 480
tcttcgggta cttcttcacg tggcggttcc ccttttaacg atctatcagg ggggaaggcc 540
ccttcgggca actcccttc cggcaactac tctcccgctc gtaccttctc accccatcc 600
acgccaacgt cccctacctc accgcttgat ttcccttctt ctcccaccaa agcagccggg 660
ggcagcacgc cggtaaccga tcatcctgac cctgttggtg gcgcgggcat cggggccgga 720
aattcgggtg ccttcaccag cgccggcgct aatcagacgg tgctgcatga caccattacc 780
gtgaaagcgg gtcaggtggt tgatggcaaa ggacaaacct tcaccgcccg ttcagaatta 840
ggcgatggcg gccagtctga aaaccagaaa ccgctgttta tactggaaga cggtgccagc 900
ctgaaaaacg tcaccatggg cgacgacggg gcggatggta ttcactctta cggtgatgcc 960
aaaatagaca atctgcacgt caccaacgtg ggtgaggacg cgattaccgt taagccaaac 1020
agcgcgggca aaaaatccca cgttgaaatc actaacagtt ccttcgagca cgcctctgac 1080
aagatcctgc agctgaatgc cgatactaac ctgagcggtg acaacgtgaa ggccaaagac 1140
tttggtactt ttgtacgcac taacggcggt caacagggta actgggatct gaatctgagc 1200
catatcagcg cagaagacgg taagttctcg ttcgttaaaa gcgatagcga ggggctaaac 1260
gtcaatacca gtgatatctc actgggtgat gttgaaaacc actacaaagt gccgatgtcc 1320
gccaacctga aggtggctga atga 1344

```

&lt;210&gt; 6

&lt;211&gt; 447

&lt;212&gt; PRT

<213> *Erwinia amylovora*

&lt;400&gt; 6

Met Ser Ile Leu Thr Leu Asn Asn Asn Thr Ser Ser Ser Pro Gly Leu

1

5

10

15



Phe Gln Ser Gly Gly Asp Asn Gly Leu Gly Gly His Asn Ala Asn Ser  
                   20                                  25                                  30

Ala Leu Gly Gln Gln Pro Ile Asp Arg Gln Thr Ile Glu Gln Met Ala  
                   35                                  40                                  45

Gln Leu Leu Ala Glu Leu Leu Lys Ser Leu Leu Ser Pro Gln Ser Gly  
                   50                                  55                                  60

Asn Ala Ala Thr Gly Ala Gly Gly Asn Asp Gln Thr Thr Gly Val Gly  
                   65                                  70                                  75                                  80

Asn Ala Gly Gly Leu Asn Gly Arg Lys Gly Thr Ala Gly Thr Thr Pro  
                                   85                                  90                                  95

Gln Ser Asp Ser Gln Asn Met Leu Ser Glu Met Gly Asn Asn Gly Leu  
                                   100                                  105                                  110

Asp Gln Ala Ile Thr Pro Asp Gly Gln Gly Gly Gly Gln Ile Gly Asp  
                   115                                  120                                  125

Asn Pro Leu Leu Lys Ala Met Leu Lys Leu Ile Ala Arg Met Met Asp  
                   130                                  135                                  140

Gly Gln Ser Asp Gln Phe Gly Gln Pro Gly Thr Gly Asn Asn Ser Ala  
                   145                                  150                                  155                                  160

Ser Ser Gly Thr Ser Ser Ser Gly Gly Ser Pro Phe Asn Asp Leu Ser  
                                   165                                  170                                  175

Gly Gly Lys Ala Pro Ser Gly Asn Ser Pro Ser Gly Asn Tyr Ser Pro  
                                   180                                  185                                  190

Val Ser Thr Phe Ser Pro Pro Ser Thr Pro Thr Ser Pro Thr Ser Pro  
                   195                                  200                                  205

---

Leu Asp Phe Pro Ser Ser Pro Thr Lys Ala Ala Gly Gly Ser Thr Pro  
                   210                                  215                                  220

Val Thr Asp His Pro Asp Pro Val Gly Ser Ala Gly Ile Gly Ala Gly  
                   225                                  230                                  235                                  240

Asn Ser Val Ala Phe Thr Ser Ala Gly Ala Asn Gln Thr Val Leu His  
                                   245                                  250                                  255

Asp Thr Ile Thr Val Lys Ala Gly Gln Val Phe Asp Gly Lys Gly Gln  
                   260                                  265                                  270





Thr Phe Thr Ala Gly Ser Glu Leu Gly Asp Gly Gly Gln Ser Glu Asn  
 275 280 285  
 Gln Lys Pro Leu Phe Ile Leu Glu Asp Gly Ala Ser Leu Lys Asn Val  
 290 295 300  
 Thr Met Gly Asp Asp Gly Ala Asp Gly Ile His Leu Tyr Gly Asp Ala  
 305 310 315 320  
 Lys Ile Asp Asn Leu His Val Thr Asn Val Gly Glu Asp Ala Ile Thr  
 325 330 335  
 Val Lys Pro Asn Ser Ala Gly Lys Lys Ser His Val Glu Ile Thr Asn  
 340 345 350  
 Ser Ser Phe Glu His Ala Ser Asp Lys Ile Leu Gln Leu Asn Ala Asp  
 355 360 365  
 Thr Asn Leu Ser Val Asp Asn Val Lys Ala Lys Asp Phe Gly Thr Phe  
 370 375 380  
 Val Arg Thr Asn Gly Gly Gln Gln Gly Asn Trp Asp Leu Asn Leu Ser  
 385 390 395 400  
 His Ile Ser Ala Glu Asp Gly Lys Phe Ser Phe Val Lys Ser Asp Ser  
 405 410 415  
 Glu Gly Leu Asn Val Asn Thr Ser Asp Ile Ser Leu Gly Asp Val Glu  
 420 425 430  
 Asn His Tyr Lys Val Pro Met Ser Ala Asn Leu Lys Val Ala Glu  
 435 440 445

&lt;210&gt; 7

&lt;211&gt; 5517

&lt;212&gt; DNA

<213> *Erwinia amylovora*

&lt;400&gt; 7

atggaattaa aatcactggg aactgaacac aaggcggcag tacacacagc ggcgcacaaac 60  
 cctgtggggc atggtgttgc cttacagcag ggcagcagca gcagcagccc gcaaaatgcc 120  
 gctgcatcat tggcggcaga aggcaaaaat cgtgggaaaa tgccgagaat tcaccagcca 180  
 tctactgcgg ctgatggtat cagcgtgtgt caccagcaaa agaaatcctt cagtctcagg 240  
 ggctgttttg ggacgaaaaa attttccaga tcggcaccgc agggccagcc aggtaccacc 300  
 cacagcaaag gggcaacatt gcgcgatctg ctggcgcggg acgacggcga aacgcagcat 360  
 gaggcggccg cgccagatgc ggcgcggttg acccggttcgg gcggcggtcaa acgccgcaat 420  
 atggacgaca tggccggggc gccaatggtg aaaggtggca gcggcgaaga taaggtacca 480



acgcagcaaa	aacggcatca	gctgaacaat	tttggccaga	tgcgccaaac	gatgttgagc	540
aaaatggctc	acccggcttc	agccaacgcc	ggcgatcgcc	tgcagcattc	accgccgcac	600
atccccggta	gccaccacga	aatcaaggaa	gaaccggttg	gctccaccag	caaggcaaca	660
acggcccacg	cagacagagt	ggaaatcgct	caggaagatg	acgacagcga	attccagcaa	720
ctgcatcaac	agcggctggc	gcgcgaacgg	gaaaatccac	cgcagccgcc	caaactcggc	780
gttgccacac	cgattagcgc	caggtttcag	cccaaactga	ctgcggttgc	ggaaagcgtc	840
cttgagggga	cagataccac	gcagtcaccc	cttaagccgc	aatcaatgct	gaaaggaagt	900
ggagccgggg	taacgccgct	ggcggtaacg	ctggataaag	gcaagttgca	gctggcaccg	960
gataatccac	ccgcgctcaa	tacgttggtg	aagcagacat	tgggtaaaga	caccagcac	1020
tatctggcgc	accatgccag	cagcgacggt	agccagcatc	tgctgctgga	caacaaaggc	1080
cacctgtttg	atatcaaaaag	caccgccacc	agctatagcg	tgctgcacaa	cagccacccc	1140
ggtgagataa	agggcaagct	ggcgcaggcg	ggtactggct	ccgtcagcgt	agacggtaaa	1200
agcggcaaga	tctcgctggg	gagcggtagc	caaagtcaca	acaaaacaat	gctaagccaa	1260
ccgggggaag	cgcaccgttc	cttattaacc	ggcatttggt	agcatcctgc	tggcgcagcg	1320
cggccgcagg	gcgagtcgat	ccgcctgcat	gacgacaaaa	ttcatatcct	gcacccggag	1380
ctgggcgtat	ggcaatctgc	ggataaagat	accacagacc	agctgtctcg	ccaggcagac	1440
ggtaagctct	atgcgctgaa	agacaaccgt	accctgcaaa	acctctccga	taataaatcc	1500
tcagaaaagc	tggctcgataa	aatcaaatcg	tattccggtg	atcagcgggg	gcagggtggcg	1560
atcctgacgg	atactcccgg	ccgccataag	atgagtatta	tgccctcgct	ggatgcttcc	1620
ccggagagcc	atatttccct	cagcctgcat	tttgccgatg	cccaccaggg	gttattgcac	1680
gggaagtcgg	agcttgaggc	acaatctgtc	gcgatcagcc	atgggagact	ggttgtggcc	1740
gatagcgaag	gcaagctggt	tagcgcgcgc	attccgaagc	aaggggatgg	aaacgaactg	1800
aaaatgaaag	ccatgcctca	gcctgcgctc	gatgaacatt	ttgggtcatga	ccaccagatt	1860
tctggatttt	tccatgacga	ccacggccag	cttaatgcgc	tgggtaaaaa	taacttcagg	1920
cagcagcatg	cctgcccgtt	gggtaacgat	catcagtttc	accccggtcg	gaacctgact	1980
gatgcgctgg	ttatcgacaa	tcagctgggg	ctgcatcata	ccaatcctga	accgcagtag	2040
attcttgata	tggggcattt	aggcagcctg	gcgttacagg	agggcaagct	tcactatttt	2100
gaccagctga	ccaaaggggtg	gactggcgcg	gagtcagatt	gtaagcagct	gaaaaaaggc	2160
ctggatggag	cagcttatct	actgaaagac	ggtgaagtga	aacgcctgaa	tattaatcag	2220
agcacctcct	ctatcaagca	cggaaacggaa	aacgtttttt	cgctgccgca	tgtgcgcaat	2280
aaaccggagc	cgggagatgc	cctgcaaggg	ctgaataaag	acgataaggc	ccaggccatg	2340
gcgggtgattg	gggtaaataa	atacctggcg	ctgacggaaa	aaggggacat	tcgctccttc	2400
cagataaaac	ccggcaccca	gcagttggag	cggccggcac	aaactctcag	ccgcgaaggc	2460
atcagcggcg	aactgaaaga	cattcatgtc	gaccacaagc	agaacctgta	tgacctgacc	2520
cacgagggag	aggtgtttca	tcagccgcgt	gaagcctggc	agaatgggtg	cgaagcagc	2580
agctggcaca	aactggcggt	gccacagagt	gaaagtaagc	taaaaagtct	ggacatgagc	2640
catgagcaca	aaccgattgc	cacctttgaa	gacggtagcc	agcatcagct	gaaggctggc	2700
ggctggcacg	cctatgcggc	acctgaacgc	gggccgctgg	cgggtgggtac	cagcggttca	2760
caaaccgtct	ttaaccgact	aatgcagggg	gtgaaaggca	aggtgatccc	aggcagcggg	2820
ttgacggtta	agctctcggc	tcagacgggg	ggaatgaccg	gcgccgaagg	gcgcaaggtc	2880
agcagtaaat	tttccgaaag	gatccgcgcc	tatgcgttca	acccaacaat	gtccacgccg	2940
cgaccgatta	aaaatgctgc	ttatgccaca	cagcacggct	ggcagggggc	tgagggggtg	3000
aagccgttgt	acgagatgca	gggagcgctg	attaaacaac	tggatgcgca	taacgttcgt	3060
cataacgcgc	cacagccaga	tttgagagc	aaactggaaa	ctctggattt	aggcgaacat	3120
ggcgcagaat	tgcttaacga	catgaagcgc	ttccgcgacg	aactggagca	gagtgcaacc	3180
cgttcgggtga	ccgttttagg	tcaacatcag	ggagtgttaa	aaagcaacgg	tgaaatcaat	3240
agcgaattta	agccatcgcc	cggcaaggcg	ttgggtccaga	gctttaacgt	caatcgctct	3300
ggtcaggatc	taagcaagtc	actgcaacag	gcagtacatg	ccacgccgcc	atccgcagag	3360



```

agtaaactgc aatccatgct ggggcacttt gtcagtgccg ggggtggatat gagtcacag 3420
aagggcgaga tcccgtggtg ccgccagcgc gatccgaatg ataaaaccgc actgaccaa 3480
tcgcgtttta ttttagatac cgtgaccatc ggtgaactgc atgaactggc cgataaggcg 3540
aaactggtat ctgaccataa acccgatgcc gatcagataa aacagctgcg ccagcagttc 3600
gatacgctgc gtgaaaagcg gtatgagagc aatccggtga agcattacac cgatatgggc 3660
ttcaccataa ataaggcgct ggaagcaaac tatgatgcgg tcaaagcctt tatcaatgcc 3720
tttaagaaag agcaccacgg cgtcaatctg accacgcgta ccgtactgga atcacagggc 3780
agtgcggagc tggcgaagaa gctcaagaat acgctgttgt ccctggacag tggtgaaagt 3840
atgagcttca gccggtcata tggcgggggc gtcagcactg tctttgtgcc tacccttagc 3900
aagaagggtg cagttccggt gatccccgga gccggcatca cgctggatcg cgcctataac 3960
ctgagcttca gtcgtaccag cggcggtatt aacgtcagtt ttggccgcga cggcggggtg 4020
agtggtaaca tcatggtcgc taccggccat gatgtgatgc cctatatgac cggtaagaaa 4080
accagtgcag gtaacgccag tgactggttg agcgcaaac ataaaatcag cccggacttg 4140
cgtatcggcg ctgctgtgag tggcacctg caaggaacgc taaaaacag cctgaagttt 4200
aagctgacag aggatgagct gcctggcttt atccatggct tgacgcattg caggttgacc 4260
ccggcagaac tggtgcaaaa ggggatcgaa catcagatga agcagggcag caaactgacg 4320
tttagcgctg atacctcgcc aaatctggat ctgcgtgccg gtatcaatct gaacgaagac 4380
ggcagtaaac caaatggtgt cactgcccgt gtttctgccg ggctaagtgc atcggaac 4440
ctggccgcgc gtcgcgtga acgcagcacc acctctggcc agtttggcag cagcacttcg 4500
gccagcaata accgccaac ctctctcaac ggggtcgccg cgggtgctaa cctgacggct 4560
gcttttagggg ttgcccattc atctacgc atcagcgcga gaagggaaac cggtcgggat cttcccgga 4620
tttacctcga ccaatgtttc ggagcgcgtg gcgctggata accgtacctc acagagtatc 4680
agcctggaat tgaagcgcgc ggagccggtg accagcaacg atatcagcga gttgacctcc 4740
acgctgggaa aacactttta ggatagcgcc acaacgaaga tgcttgccgc tctcaaagag 4800
ttagatgacg ctaagcccgc tgaacaactg catattttac agcagcattt cagtgcacaa 4860
gatgtcgtcg gtgatgaacg ctacgaggcg gtgcgcaacc tgaaaaaact ggtgatacgt 4920
caacaggctg cggacagcca cagcatggaa ttaggatctg ccagtcacag cagcactac 4980
aataatctgt cgagaataaa taatgacggc attgtcgagc tgctacacaa acatttcgat 5040
gcggcattac cagcaagcag tgccaaacgt cttggtgaaa tgatgaataa cgatccggca 5100
ctgaaagata ttattaagca gctgcaaagt acgcccgtca gcagcgccag cgtgtcgatg 5160
gagctgaaag atggtctcgc tgagcagacg gaaaaagcaa tactggacgg taaggctcgt 5220
cgtgaagaag tgggagtact tttccaggat cgtaacaact tgcgtgttaa atcggtcagc 5280
gtcagtcagt ccgtcagcaa aagcgaaggc ttcaataccc cagcgtgtt actggggacg 5340
agcaacagcg ctgctatgag catggagcgc aacatcgga ccattaattt taaatacggc 5400
caggatcaga acacccacg gcgatttacc ctggagggtg gaatagctca ggctaaccg 5460
caggtcgc atcgcgttac tgatttgaag aaggaagggc tggaaatgaa gagctaa 5517

```

&lt;210&gt; 8

&lt;211&gt; 1838

&lt;212&gt; PRT

<213> *Erwinia amylovora*

&lt;400&gt; 8

Met Glu Leu Lys Ser Leu Gly Thr Glu His Lys Ala Ala Val His Thr

1

5

10

15

Ala Ala His Asn Pro Val Gly His Gly Val Ala Leu Gln Gln Gly Ser

20

25

30



Ser Ser S r Ser Pro Gln Asn Ala Ala Ala Ser Leu Ala Ala Glu Gly  
 35 40 45  
 Lys Asn Arg Gly Lys Met Pro Arg Ile His Gln Pro Ser Thr Ala Ala  
 50 55 60  
 Asp Gly Ile Ser Ala Ala His Gln Gln Lys Lys Ser Phe Ser Leu Arg  
 65 70 75 80  
 Gly Cys Leu Gly Thr Lys Lys Phe Ser Arg Ser Ala Pro Gln Gly Gln  
 85 90 95  
 Pro Gly Thr Thr His Ser Lys Gly Ala Thr Leu Arg Asp Leu Leu Ala  
 100 105 110  
 Arg Asp Asp Gly Glu Thr Gln His Glu Ala Ala Ala Pro Asp Ala Ala  
 115 120 125  
 Arg Leu Thr Arg Ser Gly Gly Val Lys Arg Arg Asn Met Asp Asp Met  
 130 135 140  
 Ala Gly Arg Pro Met Val Lys Gly Gly Ser Gly Glu Asp Lys Val Pro  
 145 150 155 160  
 Thr Gln Gln Lys Arg His Gln Leu Asn Asn Phe Gly Gln Met Arg Gln  
 165 170 175  
 Thr Met Leu Ser Lys Met Ala His Pro Ala Ser Ala Asn Ala Gly Asp  
 180 185 190  
 Arg Leu Gln His Ser Pro Pro His Ile Pro Gly Ser His His Glu Ile  
 195 200 205  
 Lys Glu Glu Pro Val Gly Ser Thr Ser Lys Ala Thr Thr Ala His Ala  
 210 215 220  
 Asp Arg Val Glu Ile Ala Gln Glu Asp Asp Asp Ser Glu Phe Gln Gln  
 225 230 235 240  
 Leu His Gln Gln Arg Leu Ala Arg Glu Arg Glu Asn Pro Pro Gln Pro  
 245 250 255  
 Pro Lys Leu Gly Val Ala Thr Pro Ile Ser Ala Arg Phe Gln Pro Lys  
 260 265 270  
 Leu Thr Ala Val Ala Glu Ser Val Leu Glu Gly Thr Asp Thr Thr Gln  
 275 280 285





Ser Pro Leu Lys Pro Gln Ser Met Leu Lys Gly Ser Gly Ala Gly Val  
 290 295 300  
 Thr Pro Leu Ala Val Thr Leu Asp Lys Gly Lys Leu Gln Leu Ala Pro  
 305 310 315 320  
 Asp Asn Pro Pro Ala Leu Asn Thr Leu Leu Lys Gln Thr Leu Gly Lys  
 325 330 335  
 Asp Thr Gln His Tyr Leu Ala His His Ala Ser Ser Asp Gly Ser Gln  
 340 345 350  
 His Leu Leu Leu Asp Asn Lys Gly His Leu Phe Asp Ile Lys Ser Thr  
 355 360 365  
 Ala Thr Ser Tyr Ser Val Leu His Asn Ser His Pro Gly Glu Ile Lys  
 370 375 380  
 Gly Lys Leu Ala Gln Ala Gly Thr Gly Ser Val Ser Val Asp Gly Lys  
 385 390 395 400  
 Ser Gly Lys Ile Ser Leu Gly Ser Gly Thr Gln Ser His Asn Lys Thr  
 405 410 415  
 Met Leu Ser Gln Pro Gly Glu Ala His Arg Ser Leu Leu Thr Gly Ile  
 420 425 430  
 Trp Gln His Pro Ala Gly Ala Ala Arg Pro Gln Gly Glu Ser Ile Arg  
 435 440 445  
 Leu His Asp Asp Lys Ile His Ile Leu His Pro Glu Leu Gly Val Trp  
 450 455 460  
 Gln Ser Ala Asp Lys Asp Thr His Ser Gln Leu Ser Arg Gln Ala Asp  
 465 470 475 480  
 Gly Lys Leu Tyr Ala Leu Lys Asp Asn Arg Thr Leu Gln Asn Leu Ser  
 485 490 495  
 Asp Asn Lys Ser Ser Glu Lys Leu Val Asp Lys Ile Lys Ser Tyr Ser  
 500 505 510  
 Val Asp Gln Arg Gly Gln Val Ala Ile Leu Thr Asp Thr Pro Gly Arg  
 515 520 525  
 His Lys Met Ser Ile Met Pro Ser Leu Asp Ala Ser Pro Glu Ser His  
 530 535 540



Ile Ser Leu Ser Leu His Phe Ala Asp Ala His Gln Gly Leu Leu His  
 545 550 555 560

Gly Lys Ser Glu Leu Glu Ala Gln Ser Val Ala Ile Ser His Gly Arg  
 565 570 575

Leu Val Val Ala Asp Ser Glu Gly Lys Leu Phe Ser Ala Ala Ile Pro  
 580 585 590

Lys Gln Gly Asp Gly Asn Glu Leu Lys Met Lys Ala Met Pro Gln His  
 595 600 605

Ala Leu Asp Glu His Phe Gly His Asp His Gln Ile Ser Gly Phe Phe  
 610 615 620

His Asp Asp His Gly Gln Leu Asn Ala Leu Val Lys Asn Asn Phe Arg  
 625 630 635 640

Gln Gln His Ala Cys Pro Leu Gly Asn Asp His Gln Phe His Pro Gly  
 645 650 655

Trp Asn Leu Thr Asp Ala Leu Val Ile Asp Asn Gln Leu Gly Leu His  
 660 665 670

His Thr Asn Pro Glu Pro His Glu Ile Leu Asp Met Gly His Leu Gly  
 675 680 685

Ser Leu Ala Leu Gln Glu Gly Lys Leu His Tyr Phe Asp Gln Leu Thr  
 690 695 700

Lys Gly Trp Thr Gly Ala Glu Ser Asp Cys Lys Gln Leu Lys Lys Gly  
 705 710 715 720

Leu Asp Gly Ala Ala Tyr Leu Leu Lys Asp Gly Glu Val Lys Arg Leu  
 725 730 735

Asn Ile Asn Gln Ser Thr Ser Ser Ile Lys His Gly Thr Glu Asn Val  
 740 745 750

Phe Ser Leu Pro His Val Arg Asn Lys Pro Glu Pro Gly Asp Ala Leu  
 755 760 765

Gln Gly Leu Asn Lys Asp Asp Lys Ala Gln Ala Met Ala Val Ile Gly  
 770 775 780

Val Asn Lys Tyr Leu Ala Leu Thr Glu Lys Gly Asp Ile Arg Ser Phe  
 785 790 795 800



Gln Ile Lys Pro Gly Thr Gln Gln Leu Glu Arg Pro Ala Gln Thr Leu  
 805 810 815  
 Ser Arg Glu Gly Ile Ser Gly Glu Leu Lys Asp Ile His Val Asp His  
 820 825 830  
 Lys Gln Asn Leu Tyr Ala Leu Thr His Glu Gly Glu Val Phe His Gln  
 835 840 845  
 Pro Arg Glu Ala Trp Gln Asn Gly Ala Glu Ser Ser Ser Trp His Lys  
 850 855 860  
 Leu Ala Leu Pro Gln Ser Glu Ser Lys Leu Lys Ser Leu Asp Met Ser  
 865 870 875 880  
 His Glu His Lys Pro Ile Ala Thr Phe Glu Asp Gly Ser Gln His Gln  
 885 890 895  
 Leu Lys Ala Gly Gly Trp His Ala Tyr Ala Ala Pro Glu Arg Gly Pro  
 900 905 910  
 Leu Ala Val Gly Thr Ser Gly Ser Gln Thr Val Phe Asn Arg Leu Met  
 915 920 925  
 Gln Gly Val Lys Gly Lys Val Ile Pro Gly Ser Gly Leu Thr Val Lys  
 930 935 940  
 Leu Ser Ala Gln Thr Gly Gly Met Thr Gly Ala Glu Gly Arg Lys Val  
 945 950 955 960  
 Ser Ser Lys Phe Ser Glu Arg Ile Arg Ala Tyr Ala Phe Asn Pro Thr  
 965 970 975  
~~Met Ser Thr Pro Arg Pro Ile Lys Asn Ala Ala Tyr Ala Thr Gln His~~  
 980 985 990  
 Gly Trp Gln Gly Arg Glu Gly Leu Lys Pro Leu Tyr Glu Met Gln Gly  
 995 1000 1005  
 Ala Leu Ile Lys Gln Leu Asp Ala His Asn Val Arg His Asn Ala Pro  
 1010 1015 1020  
 Gln Pro Asp Leu Gln Ser Lys Leu Glu Thr Leu Asp Leu Gly Glu His  
 1025 1030 1035 1040  
 Gly Ala Glu Leu Leu Asn Asp Met Lys Arg Phe Arg Asp Glu Leu Glu  
 1045 1050 1055



Gln Ser Ala Thr Arg Ser Val Thr Val Leu Gly Gln His Gln Gly Val  
 1060 1065 1070

Leu Lys Ser Asn Gly Glu Ile Asn Ser Glu Phe Lys Pro Ser Pro Gly  
 1075 1080 1085

Lys Ala Leu Val Gln Ser Phe Asn Val Asn Arg Ser Gly Gln Asp Leu  
 1090 1095 1100

Ser Lys Ser Leu Gln Gln Ala Val His Ala Thr Pro Pro Ser Ala Glu  
 1105 1110 1115 1120

Ser Lys Leu Gln Ser Met Leu Gly His Phe Val Ser Ala Gly Val Asp  
 1125 1130 1135

Met Ser His Gln Lys Gly Glu Ile Pro Leu Gly Arg Gln Arg Asp Pro  
 1140 1145 1150

Asn Asp Lys Thr Ala Leu Thr Lys Ser Arg Leu Ile Leu Asp Thr Val  
 1155 1160 1165

Thr Ile Gly Glu Leu His Glu Leu Ala Asp Lys Ala Lys Leu Val Ser  
 1170 1175 1180

Asp His Lys Pro Asp Ala Asp Gln Ile Lys Gln Leu Arg Gln Gln Phe  
 1185 1190 1195 1200

Asp Thr Leu Arg Glu Lys Arg Tyr Glu Ser Asn Pro Val Lys His Tyr  
 1205 1210 1215

Thr Asp Met Gly Phe Thr His Asn Lys Ala Leu Glu Ala Asn Tyr Asp  
 1220 1225 1230

Ala Val Lys Ala Phe Ile Asn Ala Phe Lys Lys Glu His His Gly Val  
 1235 1240 1245

Asn Leu Thr Thr Arg Thr Val Leu Glu Ser Gln Gly Ser Ala Glu Leu  
 1250 1255 1260

Ala Lys Lys Leu Lys Asn Thr Leu Leu Ser Leu Asp Ser Gly Glu Ser  
 1265 1270 1275 1280

Met Ser Phe Ser Arg Ser Tyr Gly Gly Gly Val Ser Thr Val Phe Val  
 1285 1290 1295

Pro Thr Leu Ser Lys Lys Val Pro Val Pro Val Ile Pro Gly Ala Gly  
 1300 1305 1310





Ile Thr Leu Asp Arg Ala Tyr Asn Leu Ser Phe Ser Arg Thr Ser Gly  
 1315 1320 1325

Gly Leu Asn Val Ser Phe Gly Arg Asp Gly Gly Val Ser Gly Asn Ile  
 1330 1335 1340

Met Val Ala Thr Gly His Asp Val Met Pro Tyr Met Thr Gly Lys Lys  
 1345 1350 1355 1360

Thr Ser Ala Gly Asn Ala Ser Asp Trp Leu Ser Ala Lys His Lys Ile  
 1365 1370 1375

Ser Pro Asp Leu Arg Ile Gly Ala Ala Val Ser Gly Thr Leu Gln Gly  
 1380 1385 1390

Thr Leu Gln Asn Ser Leu Lys Phe Lys Leu Thr Glu Asp Glu Leu Pro  
 1395 1400 1405

Gly Phe Ile His Gly Leu Thr His Gly Thr Leu Thr Pro Ala Glu Leu  
 1410 1415 1420

Leu Gln Lys Gly Ile Glu His Gln Met Lys Gln Gly Ser Lys Leu Thr  
 1425 1430 1435 1440

Phe Ser Val Asp Thr Ser Ala Asn Leu Asp Leu Arg Ala Gly Ile Asn  
 1445 1450 1455

Leu Asn Glu Asp Gly Ser Lys Pro Asn Gly Val Thr Ala Arg Val Ser  
 1460 1465 1470

Ala Gly Leu Ser Ala Ser Ala Asn Leu Ala Ala Gly Ser Arg Glu Arg  
 1475 1480 1485

Ser Thr Thr Ser Gly Gln Phe Gly Ser Thr Thr Ser Ala Ser Asn Asn  
 1490 1495 1500

Arg Pro Thr Phe Leu Asn Gly Val Gly Ala Gly Ala Asn Leu Thr Ala  
 1505 1510 1515 1520

Ala Leu Gly Val Ala His Ser Ser Thr His Glu Gly Lys Pro Val Gly  
 1525 1530 1535

Ile Phe Pro Ala Phe Thr Ser Thr Asn Val Ser Ala Ala Leu Ala Leu  
 1540 1545 1550

Asp Asn Arg Thr Ser Gln Ser Ile Ser Leu Glu Leu Lys Arg Ala Glu  
 1555 1560 1565



Pro Val Thr Ser Asn Asp Ile Ser Glu Leu Thr Ser Thr Leu Gly Lys  
 1570 1575 1580

His Phe Lys Asp Ser Ala Thr Thr Lys Met Leu Ala Ala Leu Lys Glu  
 1585 1590 1595 1600

Leu Asp Asp Ala Lys Pro Ala Glu Gln Leu His Ile Leu Gln Gln His  
 1605 1610 1615

Phe Ser Ala Lys Asp Val Val Gly Asp Glu Arg Tyr Glu Ala Val Arg  
 1620 1625 1630

Asn Leu Lys Lys Leu Val Ile Arg Gln Gln Ala Ala Asp Ser His Ser  
 1635 1640 1645

Met Glu Leu Gly Ser Ala Ser His Ser Thr Thr Tyr Asn Asn Leu Ser  
 1650 1655 1660

Arg Ile Asn Asn Asp Gly Ile Val Glu Leu Leu His Lys His Phe Asp  
 1665 1670 1675 1680

Ala Ala Leu Pro Ala Ser Ser Ala Lys Arg Leu Gly Glu Met Met Asn  
 1685 1690 1695

Asn Asp Pro Ala Leu Lys Asp Ile Ile Lys Gln Leu Gln Ser Thr Pro  
 1700 1705 1710

Phe Ser Ser Ala Ser Val Ser Met Glu Leu Lys Asp Gly Leu Arg Glu  
 1715 1720 1725

Gln Thr Glu Lys Ala Ile Leu Asp Gly Lys Val Gly Arg Glu Glu Val  
 1730 1735 1740

Gly Val Leu Phe Gln Asp Arg Asn Asn Leu Arg Val Lys Ser Val Ser  
 1745 1750 1755 1760

Val Ser Gln Ser Val Ser Lys Ser Glu Gly Phe Asn Thr Pro Ala Leu  
 1765 1770 1775

Leu Leu Gly Thr Ser Asn Ser Ala Ala Met Ser Met Glu Arg Asn Ile  
 1780 1785 1790

Gly Thr Ile Asn Phe Lys Tyr Gly Gln Asp Gln Asn Thr Pro Arg Arg  
 1795 1800 1805

Phe Thr Leu Glu Gly Gly Ile Ala Gln Ala Asn Pro Gln Val Ala Ser  
 1810 1815 1820



Ala Leu Thr Asp Leu Lys Lys Glu Gly Leu Glu Met Lys Ser  
 1825 1830 1835

<210> 9  
 <211> 420  
 <212> DNA  
 <213> *Erwinia amylovora*

<400> 9  
 atgacatcgt cacagcagcg ggttgaaagg tttttacagt atttctccgc cgggtgtaaa 60  
 acgcccatac atctgaaaga cggggtgtgc gccctgtata acgaacaaga tgaggaggcg 120  
 gcgggtgctgg aagtaccgca acacagcgac agcctgttac tacactgccg aatcattgag 180  
 gctgaccac aaacttcaat aaccctgtat tcgatgctat tacagctgaa ttttgaaatg 240  
 gcggccatgc gcggctgttg gctggcgctg gatgaactgc acaacgtgcg tttatgtttt 300  
 cagcagtcgc tggagcatct ggatgaagca agttttagcg atatcgttag cggcttcac 360  
 gaacatgcgg cagaagtgcg tgagtatata gcgcaattag acgagagtag cgcggcataa 420

<210> 10  
 <211> 139  
 <212> PRT  
 <213> *Erwinia amylovora*

<400> 10  
 Met Thr Ser Ser Gln Gln Arg Val Glu Arg Phe Leu Gln Tyr Phe Ser  
 1 5 10 15

Ala Gly Cys Lys Thr Pro Ile His Leu Lys Asp Gly Val Cys Ala Leu  
 20 25 30

Tyr Asn Glu Gln Asp Glu Glu Ala Ala Val Leu Glu Val Pro Gln His  
 35 40 45

Ser Asp Ser Leu Leu Leu His Cys Arg Ile Ile Glu Ala Asp Pro Gln  
 50 55 60

Thr Ser Ile Thr Leu Tyr Ser Met Leu Leu Gln Leu Asn Phe Glu Met  
 65 70 75 80

Ala Ala Met Arg Gly Cys Trp Leu Ala Leu Asp Glu Leu His Asn Val  
 85 90 95

Arg Leu Cys Phe Gln Gln Ser Leu Glu His Leu Asp Glu Ala Ser Phe  
 100 105 110

Ser Asp Ile Val Ser Gly Phe Ile Glu His Ala Ala Glu Val Arg Glu  
 115 120 125



Tyr Ile Ala Gln Leu Asp Glu Ser Ser Ala Ala  
130 135

<210> 11

<211> 341

<212> PRT

<213> *Pseudomonas syringae*

<400> 11

Met Gln Ser Leu Ser Leu Asn Ser Ser Ser Leu Gln Thr Pro Ala Met  
1 5 10 15

Ala Leu Val Leu Val Arg Pro Glu Ala Glu Thr Thr Gly Ser Thr Ser  
20 25 30

Ser Lys Ala Leu Gln Glu Val Val Val Lys Leu Ala Glu Glu Leu Met  
35 40 45

Arg Asn Gly Gln Leu Asp Asp Ser Ser Pro Leu Gly Lys Leu Leu Ala  
50 55 60

Lys Ser Met Ala Ala Asp Gly Lys Ala Gly Gly Gly Ile Glu Asp Val  
65 70 75 80

Ile Ala Ala Leu Asp Lys Leu Ile His Glu Lys Leu Gly Asp Asn Phe  
85 90 95

Gly Ala Ser Ala Asp Ser Ala Ser Gly Thr Gly Gln Gln Asp Leu Met  
100 105 110

Thr Gln Val Leu Asn Gly Leu Ala Lys Ser Met Leu Asp Asp Leu Leu  
115 120 125

Thr Lys Gln Asp Gly Gly Thr Ser Phe Ser Glu Asp Asp Met Pro Met  
130 135 140

Leu Asn Lys Ile Ala Gln Phe Met Asp Asp Asn Pro Ala Gln Phe Pro  
145 150 155 160

Lys Pro Asp Ser Gly Ser Trp Val Asn Glu Leu Lys Glu Asp Asn Phe  
165 170 175

Leu Asp Gly Asp Glu Thr Ala Ala Phe Arg Ser Ala Leu Asp Ile Ile  
180 185 190

Gly Gln Gln Leu Gly Asn Gln Gln Ser Asp Ala Gly Ser Leu Ala Gly





<400> 12						
atgcagagtc	tcagtcttaa	cagcagctcg	ctgcaaacc	cggcaatggc	ccttgtcctg	60
gtacgtcctg	aagccgagac	gactggcagt	acgtcgagca	aggcgcttca	ggaagttgtc	120
gtgaagctgg	ccgaggaact	gatgcgcaat	ggткаactcg	acgacagctc	gccattggga	180
aaactgttgg	ccaagtcgat	ggccgcagat	ggcaaggcgg	gcgggcggat	tgaggatgtc	240
atcgctgcgc	tggacaagct	gatccatgaa	aagctcggtg	acaacttcgg	cgcgtctgcg	300
gacagcgcct	cgggtaccgg	acagcaggac	ctgatgactc	agggtgctcaa	tggcctggcc	360
aagtcgatgc	tcgatgatct	tctgaccaag	caggatggcg	ggacaagctt	ctccgaagac	420
gatatgccga	tgctgaacaa	gatcgcgcag	ttcatggatg	acaatcccgc	acagtttccc	480
aagccggact	cgggctcctg	ggtgaacgaa	ctcaaggaag	acaacttcct	tgatggcgac	540
gaaacggctg	cgttccgttc	ggcactcgac	atcattggcc	agcaactggg	taatcagcag	600
agtgacgctg	gcagtctggc	agggacgggt	ggaggtctgg	gcactccgag	cagtttttcc	660
aacaactcgt	ccgtgatggg	tgatccgctg	atcgacgcc	ataccgggtcc	cggtgacagc	720



ggcaataccc gtggtgaagc ggggcaactg atcggcgagc ttatcgaccg tggcctgcaa 780  
 tccgtattgg ccggtggtgg actgggcaca cccgtaaaca ccccgagac cggtagctcg 840  
 gcgaatggcg gacagtccgc tcaggatctt gatcagttgc tggcgggctt gctgctcaag 900  
 ggcctggagg caacgctcaa ggatgccggg caaacaggca ccgacgtgca gtcgagcgct 960  
 gcgcaaactc ccaccttgct ggtcagtagc ctgctgcaag gcacccgcaa tcaggctgca 1020  
 gcctga 1026

&lt;210&gt; 13

&lt;211&gt; 1729

&lt;212&gt; DNA

<213> *Pseudomonas syringae*

&lt;400&gt; 13

tccacttcgc tgattttgaa attggcagat tcatagaaac gttcaggtgt ggaaatcagg 60  
 ctgagtgcgc agatttcgtt gataaggggtg tggtagctgg cattgttggt catttcaagg 120  
 cctctgagtg cgggtgcggag caataccagt cttcctgctg gcgtgtgcac actgagtcgc 180  
 aggcataaggc atttcagttc cttgcgttgg ttgggcatat aaaaaaagga acttttaaaa 240  
 acagtgcaat gagatgccgg caaaacggga accggctcgt gcgctttgcc actcacttcg 300  
 agcaagctca accccaaaca tccacatccc tatcgaacgg acagcgatac ggccacttgc 360  
 tctggtaaac cctggagctg gcgtcgggtcc aattgccac ttagcgaggt aacgcagcat 420  
 gagcatcggc atcacacccc ggccgcaaca gaccaccacg ccactcgatt tttcggcgct 480  
 aagcggcaag agtcctcaac caaacacgtt cggcgagcag aacactcagc aagcgatcga 540  
 cccgagtgcg ctgttggttcg gcagcgacac acagaaagac gtcaacttcg gcacgcccga 600  
 cagcaccgtc cagaatccgc aggacgccag caagcccaac gacagccagt ccaacatcgc 660  
 taaattgatc agtgattga tcatgtcgtt gctgcagatg ctcaccaact ccaataaaaa 720  
 gcaggacacc aatcaggaac agcctgatag ccaggtcctt ttccagaaca acggcgggct 780  
 cggtagaccg tcggccgata gcggggggcg cggtagaccg gatgcgacag gtggcggcg 840  
 cggtgatacg ccaagcgcaa caggcgggtg cggcgggtgat actccgaccg caacaggcgg 900  
 tggcggcagc ggtggcggcg gcacacccac tgcaacaggt ggccggcagcg gtggcacacc 960  
 cactgcaaca ggcggtggcg aggggtggcgt aacaccgcaa atcactccgc agttggccaa 1020  
 ccctaaccgt acctcaggta ctggctcggg gtccggacacc gcagggttcta ccgagcaagc 1080  
 cggcaagatc aatgtggtga aagacaccat caaggtcggc gctggcggaag tctttgacgg 1140  
 ccacggcgca accttcactg ccgacaaatc tatgggtaac ggagaccagg gcgaaaatca 1200  
 gaagcccatg ttcgagctgg ctgaaggcgc tacgttgaag aatgtgaacc tgggtgagaa 1260  
 cgaggtcgat ggcattccacg tgaaagccaa aaacgctcag gaagtcacca ttgacaacgt 1320  
 gcatgccagc aacgtcgggtg aagacctgat tacggtcaaa ggccagggag gcgcagcggt 1380  
 cactaatctg aacatcaaga acagcagtg caaagggtgca gacgacaagg ttgtccagct 1440  
 caacgccaac actcacttga aaatcgacaa cttcaaggcc gacgatttcg gcacgatggg 1500  
 tcgcaccaac ggtggcaagc agtttgatga catgagcatc gagctgaacg gcatcgaagc 1560  
 taaccacggc aagttcgccc tggtgaaaag cgacagtgac gatctgaagc tggcaacggg 1620  
 caacatcgcc atgaccgacg tcaaacacgc ctacgataaa acccaggcat cgaccaaca 1680  
 caccgagctt tgaatccaga caagtagctt gaaaaaaggg ggtggactc 1729

&lt;210&gt; 14

&lt;211&gt; 424

&lt;212&gt; PRT

<213> *Pseudomonas syringae*



&lt;400&gt; 14

Met Ser Ile Gly Ile Thr Pro Arg Pro Gln Gln Thr Thr Thr Pro Leu  
 1 5 10 15

Asp Phe Ser Ala Leu Ser Gly Lys Ser Pro Gln Pro Asn Thr Phe Gly  
 20 25 30

Glu Gln Asn Thr Gln Gln Ala Ile Asp Pro Ser Ala Leu Leu Phe Gly  
 35 40 45

Ser Asp Thr Gln Lys Asp Val Asn Phe Gly Thr Pro Asp Ser Thr Val  
 50 55 60

Gln Asn Pro Gln Asp Ala Ser Lys Pro Asn Asp Ser Gln Ser Asn Ile  
 65 70 75 80

Ala Lys Leu Ile Ser Ala Leu Ile Met Ser Leu Leu Gln Met Leu Thr  
 85 90 95

Asn Ser Asn Lys Lys Gln Asp Thr Asn Gln Glu Gln Pro Asp Ser Gln  
 100 105 110

Ala Pro Phe Gln Asn Asn Gly Gly Leu Gly Thr Pro Ser Ala Asp Ser  
 115 120 125

Gly Gly Gly Gly Thr Pro Asp Ala Thr Gly Gly Gly Gly Gly Asp Thr  
 130 135 140

Pro Ser Ala Thr Gly Gly Gly Gly Gly Asp Thr Pro Thr Ala Thr Gly  
 145 150 155 160

Gly Gly Gly Ser Gly Gly Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly  
 165 170 175

Ser Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly Glu Gly Gly Val Thr  
 180 185 190

Pro Gln Ile Thr Pro Gln Leu Ala Asn Pro Asn Arg Thr Ser Gly Thr  
 195 200 205

Gly Ser Val Ser Asp Thr Ala Gly Ser Thr Glu Gln Ala Gly Lys Ile  
 210 215 220

Asn Val Val Lys Asp Thr Ile Lys Val Gly Ala Gly Glu Val Phe Asp  
 225 230 235 240

Gly His Gly Ala Thr Phe Thr Ala Asp Lys Ser Met Gly Asn Gly Asp  
 245 250 255



Gln Gly Glu Asn Gln Lys Pr Met Ph Glu Leu Ala Glu Gly Ala Thr  
 260 265 270  
 Leu Lys Asn Val Asn Leu Gly Glu Asn Glu Val Asp Gly Ile His Val  
 275 280 285  
 Lys Ala Lys Asn Ala Gln Glu Val Thr Ile Asp Asn Val His Ala Gln  
 290 295 300  
 Asn Val Gly Glu Asp Leu Ile Thr Val Lys Gly Glu Gly Gly Ala Ala  
 305 310 315 320  
 Val Thr Asn Leu Asn Ile Lys Asn Ser Ser Ala Lys Gly Ala Asp Asp  
 325 330 335  
 Lys Val Val Gln Leu Asn Ala Asn Thr His Leu Lys Ile Asp Asn Phe  
 340 345 350  
 Lys Ala Asp Asp Phe Gly Thr Met Val Arg Thr Asn Gly Gly Lys Gln  
 355 360 365  
 Phe Asp Asp Met Ser Ile Glu Leu Asn Gly Ile Glu Ala Asn His Gly  
 370 375 380  
 Lys Phe Ala Leu Val Lys Ser Asp Ser Asp Asp Leu Lys Leu Ala Thr  
 385 390 395 400  
 Gly Asn Ile Ala Met Thr Asp Val Lys His Ala Tyr Asp Lys Thr Gln  
 405 410 415  
 Ala Ser Thr Gln His Thr Glu Leu  
 420

&lt;210&gt; 15

&lt;211&gt; 344

&lt;212&gt; PRT

&lt;213&gt; Pseudomonas solanacearum

&lt;400&gt; 15

Met Ser Val Gly Asn Ile Gln Ser Pro Ser Asn Leu Pro Gly Leu Gln  
 1 5 10 15  
 Asn Leu Asn Leu Asn Thr Asn Thr Asn Ser Gln Gln Ser Gly Gln Ser  
 20 25 30  
 Val Gln Asp Leu Ile Lys Gln Val Glu Lys Asp Ile Leu Asn Ile Ile





35	40	45
Ala Ala Leu Val Gln Lys Ala Ala Gln Ser Ala Gly Gly Asn Thr Gly		
50	55	60
Asn Thr Gly Asn Ala Pro Ala Lys Asp Gly Asn Ala Asn Ala Gly Ala		
65	70	75
80		
Asn Asp Pro Ser Lys Asn Asp Pro Ser Lys Ser Gln Ala Pro Gln Ser		
85	90	95
100		
Ala Asn Lys Thr Gly Asn Val Asp Asp Ala Asn Asn Gln Asp Pro Met		
105	110	
Gln Ala Leu Met Gln Leu Leu Glu Asp Leu Val Lys Leu Leu Lys Ala		
115	120	125
Ala Leu His Met Gln Gln Pro Gly Gly Asn Asp Lys Gly Asn Gly Val		
130	135	140
Gly Gly Ala Asn Gly Ala Lys Gly Ala Gly Gly Gln Gly Gly Leu Ala		
145	150	155
160		
Glu Ala Leu Gln Glu Ile Glu Gln Ile Leu Ala Gln Leu Gly Gly Gly		
165	170	175
Gly Ala Gly Ala Gly Gly Ala Gly Gly Gly Val Gly Gly Ala Gly Gly		
180	185	190
Ala Asp Gly Gly Ser Gly Ala Gly Gly Ala Gly Gly Ala Asn Gly Ala		
195	200	205
Asp Gly Gly Asn Gly Val Asn Gly Asn Gln Ala Asn Gly Pro Gln Asn		
210	215	220
Ala Gly Asp Val Asn Gly Ala Asn Gly Ala Asp Asp Gly Ser Glu Asp		
225	230	235
240		
Gln Gly Gly Leu Thr Gly Val Leu Gln Lys Leu Met Lys Ile Leu Asn		
245	250	255
Ala Leu Val Gln Met Met Gln Gln Gly Gly Leu Gly Gly Gly Asn Gln		
260	265	270
Ala Gln Gly Gly Ser Lys Gly Ala Gly Asn Ala Ser Pro Ala Ser Gly		
275	280	285
Ala Asn Pro Gly Ala Asn Gln Pro Gly Ser Ala Asp Asp Gln Ser Ser		



290

295

300

Gly Gln Asn Asn Leu Gln Ser Gln Ile Met Asp Val Val Lys Glu Val  
 305 310 315 320

Val Gln Ile Leu Gln Gln Met Leu Ala Ala Gln Asn Gly Gly Ser Gln  
 325 330 335

Gln Ser Thr Ser Thr Gln Pro Met  
 340

&lt;210&gt; 16

&lt;211&gt; 1035

&lt;212&gt; DNA

<213> *Pseudomonas solanacearum*

&lt;400&gt; 16

atgtcagtcg gaaacatcca gagcccgctcg aacctcccgg gtctgcagaa cctgaacctc 60  
 aacaccaaca ccaacagcca gcaatcgggc cagtcctgtc aagacctgat caagcaggtc 120  
 gagaaggaca tcctcaacat catcgcagcc ctctgtcaga aggccgcaca gtcggcgggc 180  
 ggcaacaccg gtaacaccgg caacgcgccg gcgaaggacg gcaatgccaa cgcggggcgcc 240  
 aacgacccga gcaagaacga cccgagcaag agccaggctc cgcagtcggc caacaagacc 300  
 ggcaacgtcg acgacgccaa caaccaggat ccgatgcaag cgctgatgca gctgctggaa 360  
 gacctggtga agctgctgaa ggcgggcctg cacatgcagc agccccggcg caatgacaag 420  
 ggcaacggcg tggcggtgc caacggcgcc aagggtgccc gcggccaggc cggcctggcc 480  
 gaagcgctgc aggagatcga gcagatcctc gccagctcg gcggcgcgcg tgctggcgcc 540  
 ggcgggcgcg gtggcggtgt cggcggtgct ggtggcgcg atggcggtc cggcgcggt 600  
 ggcgcgaggc gtgcgaacgg cgccgacggc ggcaatggcg tgaacggcaa ccaggcgaac 660  
 ggccccgaga acgcaggcga tgtcaacggt gccaacggcg cggatgacgg cagcgaagac 720  
 caggcgcgcc tcaccggcgt gctgcaaaag ctgatgaaga tcctgaacgc gctggtgcag 780  
 atgatgcagc aaggcggcct cggcgggcgc aaccaggcgc agggcggtc gaagggtgcc 840  
 ggcaacgcct cgccggcttc cggcgcgaa cggggcgcg accagcccgg ttcggcggt 900  
 gatcaatcgt ccggccagaa caatctgcaa tcccagatca tggatgtggt gaaggaggtc 960  
 gtccagatcc tgcagcagat gctggcgggc cagaacggcg gcagccagca gtccacctc 1020  
 acgcagccga tgtaa 1035

&lt;210&gt; 17

&lt;211&gt; 26

&lt;212&gt; PRT

<213> *Xanthomonas campestris* pv. *glycines*

&lt;400&gt; 17

Thr Leu Ile Glu Leu Met Ile Val Val Ala Ile Ile Ala Ile Leu Ala  
 1 5 10 15

Ala Ile Ala Leu Pro Ala Tyr Gln Asp Tyr  
 20 25



&lt;210&gt; 18

&lt;211&gt; 20

&lt;212&gt; PRT

<213> *Xanthomonas campestris* pv. *pelargonii*

&lt;400&gt; 18

Ser Ser Gln Gln Ser Pro Ser Ala Gly Ser Glu Gln Gln Leu Asp Gln

1

5

10

15

Leu Leu Ala Met

20



# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/26039

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/82 C12N15/31 A01N63/02

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 32844 A (CORNELL RES FOUNDATION INC) 30 July 1998 (1998-07-30) claims	1-30
A	WO 96 39802 A (CORNELL RES FOUNDATION INC) 19 December 1996 (1996-12-19) claims	1-30
A	WO 98 24297 A (CORNELL RES FOUNDATION INC) 11 June 1998 (1998-06-11) claims	1-30
A	WO 98 37752 A (CORNELL RES FOUNDATION INC) 3 September 1998 (1998-09-03) claims	1-30

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 May 2000

Date of mailing of the international search report

07/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Decorte, D

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 99/26039

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9832844 A	30-07-1998	AU 6043198 A PL 334907 A	18-08-1998 27-03-2000
WO 9639802 A	19-12-1996	US 5650387 A AU 714512 B AU 5982196 A BR 9609073 A CA 2223616 A CN 1192647 A EP 0871354 A FI 974430 A JP 11506938 T NZ 309611 A PL 323823 A US 5859324 A US 5776889 A	22-07-1997 06-01-2000 30-12-1996 26-01-1999 19-12-1996 09-09-1998 21-10-1998 26-01-1998 22-06-1999 30-08-1999 27-04-1998 12-01-1999 07-07-1998
WO 9824297 A	11-06-1998	AU 5693598 A CN 1245393 A EP 0957672 A PL 334126 A	29-06-1998 23-02-2000 24-11-1999 14-02-2000
WO 9837752 A	03-09-1998	AU 6666498 A BR 9807632 A US 5977060 A	18-09-1998 22-02-2000 02-11-1999